



US009951129B2

(12) **United States Patent**
Towne et al.

(10) **Patent No.:** **US 9,951,129 B2**
(45) **Date of Patent:** **Apr. 24, 2018**

(54) **HUMAN IL-23 ANTIGEN BINDING PROTEINS**

(71) Applicant: **AMGEN INC.**, Thousand Oaks, CA (US)

(72) Inventors: **Jennifer E. Towne**, Seattle, WA (US); **Janet D. Cheng**, Seattle, WA (US); **Jason C. O'Neill**, Brier, WA (US); **Yu Zhang**, Shoreline, WA (US); **Yu Sun**, Seattle, WA (US); **Heather Cerne**, Seattle, WA (US); **Derek E. Piper**, Santa Clara, CA (US); **Randal R. Ketchum**, Snohomish, WA (US)

(73) Assignee: **AMGEN INC.**, Thousand Oaks, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/271,315**

(22) Filed: **Sep. 21, 2016**

(65) **Prior Publication Data**

US 2017/0002070 A1 Jan. 5, 2017

Related U.S. Application Data

(60) Division of application No. 14/228,556, filed on Mar. 28, 2014, now Pat. No. 9,487,580, which is a continuation of application No. 13/504,449, filed as application No. PCT/US2010/054148 on Oct. 26, 2010, now Pat. No. 8,722,033.

(60) Provisional application No. 61/381,287, filed on Sep. 9, 2010, provisional application No. 61/254,982, filed on Oct. 26, 2009.

(51) **Int. Cl.**
C12N 15/24 (2006.01)
C07K 16/24 (2006.01)
A61K 39/395 (2006.01)
A61K 47/68 (2017.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
CPC **C07K 16/244** (2013.01); **A61K 39/395** (2013.01); **A61K 47/6845** (2017.08); **C07K 16/24** (2013.01); **A61K 2039/505** (2013.01); **C07K 2317/21** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/31** (2013.01); **C07K 2317/54** (2013.01); **C07K 2317/55** (2013.01); **C07K 2317/565** (2013.01); **C07K 2317/622** (2013.01); **C07K 2317/626** (2013.01); **C07K 2317/76** (2013.01); **C07K 2317/92** (2013.01)

(58) **Field of Classification Search**
CPC **C07K 16/244**; **C12N 15/00**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,399,216 A	8/1983	Axel et al.
4,740,461 A	4/1988	Kaufman
4,751,180 A	6/1988	Cousens et al.
4,816,567 A	3/1989	Cabilly et al.
4,912,040 A	3/1990	Kaufman et al.
4,935,233 A	6/1990	Bell et al.
4,946,778 A	8/1990	Ladner et al.
4,959,455 A	9/1990	Clark et al.
4,965,195 A	10/1990	Namen et al.
4,968,607 A	11/1990	Dower et al.
5,011,912 A	4/1991	Hopp et al.
5,260,203 A	11/1993	Ladner et al.
5,262,522 A	11/1993	Gearing
5,426,048 A	6/1995	Gearing
5,457,035 A	10/1995	Baum et al.
5,457,038 A	10/1995	Trinchieri et al.
5,530,101 A	6/1996	Queen et al.
5,545,806 A	8/1996	Lonberg et al.
5,545,807 A	8/1996	Surani et al.
5,569,825 A	10/1996	Lonberg et al.
5,585,089 A	12/1996	Queen et al.
5,625,126 A	4/1997	Lonberg et al.
5,633,425 A	5/1997	Lonberg et al.
5,661,016 A	8/1997	Lonberg et al.
5,693,761 A	12/1997	Queen et al.
5,693,762 A	12/1997	Queen et al.
5,770,429 A	6/1998	Lonberg et al.
5,780,597 A	7/1998	Gately et al.
5,789,650 A	8/1998	Lonberg et al.
5,814,318 A	9/1998	Lonberg et al.
5,874,299 A	2/1999	Lonberg et al.
5,877,397 A	3/1999	Lonberg et al.
6,162,963 A	12/2000	Kucherlapati et al.
6,171,586 B1	1/2001	Lam et al.
6,180,370 B1	1/2001	Queen et al.
6,255,458 B1	7/2001	Lonberg et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU	2010291927 A1	4/2012
EP	0367566 A1	5/1990

(Continued)

OTHER PUBLICATIONS

The CCP4 Suite: Programs for Protein Crystallography, *Acta Crystallogr D Biol Crystallogr*, 50(Pt 5): p. 760-3 (1994).
Adams et al., The c-myc oncogene driven by immunoglobulin enhancers induces lymphoid malignancy in transgenic mice, *Nature*, 318:533-538 (1985).
Afzali et al., The role of T helper 17 (Th17) and regulatory T cells (Treg) in human organ transplantation and autoimmune disease, *Clinical and Experimental Immunology*, 148:32-46 (2007).

(Continued)

Primary Examiner — Prema Mertz

(74) *Attorney, Agent, or Firm* — Marshall, Gerstein & Borun LLP

(57) **ABSTRACT**

Antigen binding proteins that bind to human IL-23 protein are provided. Nucleic acids encoding the antigen binding protein, vectors, and cells encoding the same as well as use of IL-23 antigen binding proteins for diagnostic and therapeutic purposes are also provided.

18 Claims, 2 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

6,270,964	B1	8/2001	Michnick et al.
6,300,129	B1	10/2001	Lonberg et al.
6,495,667	B1	12/2002	Bazan
6,673,986	B1	1/2004	Kucherlapati et al.
6,713,610	B1	3/2004	Kucherlapati et al.
6,914,128	B1	7/2005	Salfeld et al.
7,090,847	B1	8/2006	Oppmann et al.
7,247,711	B2	7/2007	Benson et al.
7,282,204	B2	10/2007	Ofit et al.
7,491,391	B2	2/2009	Benson et al.
7,501,247	B2	3/2009	Kastelein et al.
7,504,485	B2	3/2009	Salfeld et al.
7,510,709	B2	3/2009	Gurney
7,575,741	B2	8/2009	Bowman et al.
7,608,690	B2	10/2009	Bazan
7,700,739	B2	4/2010	Lacy et al.
7,740,848	B2	6/2010	Kastelein et al.
7,750,126	B2	7/2010	Hirata
7,790,862	B2	9/2010	Lewis et al.
7,807,160	B2	10/2010	Presta et al.
7,807,414	B2	10/2010	Benson et al.
7,820,168	B2	10/2010	Cua et al.
7,872,102	B2	1/2011	Beidler et al.
7,883,704	B2	2/2011	Salfeld et al.
7,893,215	B2	2/2011	Bowman et al.
7,910,703	B2	3/2011	Lewis et al.
7,935,344	B2	5/2011	Benson et al.
7,993,645	B2	8/2011	Benson et al.
8,080,247	B2	12/2011	Giles-Komar et al.
8,106,177	B2	1/2012	Benson et al.
8,110,187	B2	2/2012	Gately et al.
8,182,810	B2	5/2012	Bazan
8,221,760	B2	7/2012	Benson et al.
8,227,579	B2	7/2012	Lewis et al.
8,287,869	B2	10/2012	Gurney
8,293,883	B2	10/2012	Presta
8,329,170	B2	12/2012	Giles-Komar et al.
8,333,968	B2	12/2012	Lewis et al.
8,362,212	B2	1/2013	Presta
8,404,813	B2	3/2013	Presta
8,563,697	B2	10/2013	Clarke et al.
8,586,035	B2	11/2013	Kopf et al.
8,778,346	B2	7/2014	Barrett et al.
8,927,693	B2	1/2015	Dasgupta et al.
2007/0048315	A1	3/2007	Presta
2007/0218064	A1	9/2007	Benson et al.
2009/0123479	A1	5/2009	Bembridge et al.
2010/0143357	A1	6/2010	Cua et al.
2011/0177022	A1	7/2011	Oppmann et al.
2011/0262445	A1	10/2011	Clarke et al.
2011/0319292	A1	12/2011	Benson et al.
2012/0027799	A1	2/2012	Sears et al.
2012/0128689	A1	5/2012	Clarkson et al.
2012/0183964	A1	7/2012	Bazan
2012/0264917	A1	10/2012	Saunders et al.
2012/0276105	A1	11/2012	Kastelein et al.
2012/0282269	A1	11/2012	Barrett et al.
2012/0308573	A1	12/2012	Benson et al.
2013/0004501	A1	1/2013	Towne et al.
2013/0039916	A1	2/2013	Presta et al.
2013/0122009	A1	5/2013	Presta et al.

FOREIGN PATENT DOCUMENTS

EP	0460846	A1	12/1991
EP	546073	A1	6/1993
EP	546073	B1	9/1997
EP	1210434	B1	6/2002
EP	1002084	B1	9/2009
EP	1072610	B1	7/2010
WO	WO-1987/005330	A1	9/1987
WO	WO-1988/001649	A1	3/1988
WO	WO-1990/004036	A1	4/1990
WO	WO-1991/010741	A1	7/1991
WO	WO-1992/003918	A1	3/1992

WO	WO-1992/022646	A1	12/1992
WO	WO-1993/001227	A1	1/1993
WO	WO-1993/010151	A1	5/1993
WO	WO-1994/002602	A1	2/1994
WO	WO-1994/010308	A1	5/1994
WO	WO-1996/033735	A1	10/1996
WO	WO-1998/024893	A2	6/1998
WO	WO-1999/005280	A1	2/1999
WO	WO-1999/010494	A2	3/1999
WO	WO-2000/009560	A2	2/2000
WO	WO-2004/058178	A2	7/2004
WO	WO-2004/081190	A2	9/2004
WO	WO-2004/101750	A2	11/2004
WO	WO-2006/036745	A2	4/2006
WO	WO-2006/068987	A2	6/2006
WO	WO-2006/069036	A2	6/2006
WO	WO-2007/005955	A2	1/2007
WO	WO-2007/024846	A2	3/2007
WO	WO-2007/027714	A2	3/2007
WO	WO-2007/076523	A2	7/2007
WO	WO-2007/076524	A2	7/2007
WO	WO-2007/147019	A2	12/2007
WO	WO-2008/088823	A2	7/2008
WO	WO-2008/103432	A1	8/2008
WO	WO-2008/103473	A1	8/2008
WO	WO-2008/153610	A2	12/2008
WO	WO-2009/043933	A1	4/2009
WO	WO-2009/068627	A2	6/2009
WO	WO-2009/082624	A2	7/2009
WO	WO-2009/117289	A2	9/2009
WO	WO-2010/017598	A1	2/2010
WO	WO-2010/027766	A1	3/2010
WO	WO-2010/115786	A1	10/2010
WO	WO-2010/142534	A1	12/2010
WO	WO-2011/032148	A1	3/2011
WO	WO-2011/056600	A1	5/2011
WO	WO-2011/088120	A1	7/2011
WO	WO-2011/103105	A1	8/2011
WO	WO-2012/009760	A1	1/2012
WO	WO-2012/045848	A1	4/2012
WO	WO-2012/048134	A2	4/2012
WO	WO-2012/061448	A1	5/2012
WO	WO-2012/094623	A2	7/2012

OTHER PUBLICATIONS

Ahern et al., The interleukin-23 axis in intestinal inflammation, *Immun. Rev.*, 226:147-159 (2008).

Alexander et al., Expression of the c-myc Oncogene under control of an immunoglobulin enhancer in Eμ-myc Transgenic Mice, *Mol. Cell. Biol.*, 7:1436-1444 (1987).

Amit et al., Three-dimensional structure of an antigen-antibody complex at 2.8 Å resolution, *Science*, 233(4765):747-53 (1986).

Aplin and Wriston, Preparation, Properties, and Applications of Carbohydrate Conjugates of Proteins and Lipids, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

Ashkenazi et al., Protection against endotoxin shock by a tumor necrosis factor receptor immunoadhesin, *Proc. Natl. Acad. Sci. USA*, 88:10535 (1991).

Baum et al., Identification of ligands for OX40 receptor, *EMBO J.*, 13:3992-4001 (1994).

Belladonna et al., IL-23 and IL-12 have overlapping, but distinct, effects on murine dendritic cells, *The Journal of Immunology*, 168:5448-5454 (2002).

Benoist and Chambon, in vivo sequence requirements of the SV40 early promoter region, *Nature*, 290:304-310 (1981).

Beyer et al., Crystal structures of the pro-inflammatory cytokine interleukin-23 and its complex with a high-affinity neutralizing antibody, *J. Mol. Biol.*, 382(4):942-55 (2008).

Beyer, et al., Crystal Structures of the Pro-Inflammatory Cytokine Interleukin-23 and Its Complex with a High-Affinity Neutralizing Antibody, *J Mol Biol*, 382(4):942-55 (2008).

Bianchi and McGrew, High-Level Expression of Full-Length Antibodies Using Trans-Complementing Expression Vectors, *Biotech. Bioeng.*, 84:439-44 (2003).

Bird et al., Single-Chain Antigen-Binding Proteins, *Science*, 242:423 (1988).

(56)

References Cited

OTHER PUBLICATIONS

- Blauvelt, in this Issue—Full Court Press on Psoriasis, *The Journal of Investigative Dermatology*, pp. vii-viii (2004).
- Bloom et al., Intrachain disulfide bond in the core hinge region of human IgG4, *Protein Science*, 6:407 (1997).
- Bowie et al., A method to identify protein sequences that fold into a known three-dimensional structure, *Science*, 253:164-170 (1991).
- Bowman et al., Rationale and safety of anti-interleukin-23 and anti-interleukin-17A therapy, *Current Opinion in Infectious Diseases*, 19:245-52 (2006).
- Brenner et al., Population statistics of protein structures: lessons from structural classifications, *Curr. Op. Struct. Biol.*, 7:369-376 (1997).
- Brinster et al., Regulation of metallothionein—thymidine kinase fusion plasmids injected into mouse eggs, *Nature*, 296:39-42 (1982).
- Brown et al., Tolerance of single, but not multiple, amino acid replacements in antibody VH CDR 2: a means of minimizing B cell wastage from somatic hypermutation?, *J. Immunol.*, 156(9):3285-91 (1996).
- Bruggermann et al., Designer mice: the production of human antibody repertoires in transgenic animals, *Year in Immunol.*, 7:33-40 (1993).
- Brunger, et al., Crystallography & NMR System: A new software suite for macromolecular structure determination, *Acta Crystallogr. D Biol. Crystallogr.*, 54(Pt 5): p. 905-21 (1998).
- Byrn et al., Biological properties of a CD4 immunoadhesin, *Nature*, 344:677 (1990).
- Carillo et al., The Multiple Sequence Alignment Problem in Biology, *SIAM J. Applied Math.*, 48:1073 (1988).
- Chen et al., Anti-IL-23 therapy inhibits multiple inflammatory pathways and ameliorates autoimmune encephalomyelitis, *The Journal of Clinical Investigation*, 116:1317-1326 (2006).
- Chen et al., Immunoglobulin gene rearrangement in B cell deficient mice generated by targeted deletion of the JH locus, *International Immunology*, 5:647-656 (1993).
- Cheung, et al., Epitope-Specific Antibody Response to the Surface Antigen of Duck Hepatitis B Virus in Infected Ducks, *Virology*, 176:546-552 (1990).
- Chothia & Lesk, Canonical Structures for the Hypervariable Regions of Immunoglobulins, *J. Mol. Biol.*, 196:901-917 (1987).
- Chothia et al., Conformations of immunoglobulin hypervariable regions, *Nature*, 342:878-883 (1989).
- Chou et al., Conformational Parameters for Amino Acids in Helical, β -Sheet, and Random Coil Regions Calculated from Proteins, *Biochemistry*, 113:211-222 (1974).
- Chou et al., Empirical Predictions of Proteins Conformation, *Ann. Rev. Biochem.*, 47:251-276 (1978).
- Chou et al., Prediction of Protein Conformation, *Biochem.*, 13:222-245 (1974).
- Chou et al., Prediction of the secondary structure of proteins from their amino acid sequence, *Adv. Enzymol. Relat. Areas Mol. Biol.*, 47:45-148 (1978).
- Chou et al., Prediction of β -Turns, *Biophys. J.*, 26:367-384 (1979).
- ClinicalTrials.gov, Identifier: NCT01094093, Safety, Tolerability, Pharmacokinetics and Pharmacodynamics of AMG 139 in Healthy Subjects and Subjects With Moderate to Severe Psoriasis, dated Mar. 25, 2010.
- Cosman et al., Cloning, sequence and expression of human interleukin-2 receptor, *Nature*, 312:768 (1984).
- Cua et al., Interleukin-23 rather than Interleukin-12 is the critical cytokine for autoimmune inflammation of the brain, *Nature*, 421:744-748 (2003).
- Davies et al., Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding, *Immunotechnology*, 2(3):169-79 (1996).
- Dayhoff et al., A model of evolutionary change in proteins, *Atlas of Protein Sequence and Structure*, 5:345-352 (1978).
- de Graaf et al., Expression of scFvs and scFv Fusion Proteins in Eukaryotic Cells, *Methods Mol Biol.*, 178:379-387 (2002).
- DeBoer et al., The tac promoter: A functional hybrid derived from the trp and lac promoters, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25 (1983).
- Devereux et al., A comprehensive set of sequence analysis programs for the VAX, *Nucl. Acid Res.*, 12:387 (1984).
- Duskin et al., Relationship of the structure and biological activity of the natural homologues of tunicamycin, *J. Biol. Chem.*, 257:3105 (1982).
- Edge et al., Deglycosylation of Glycoproteins by Trifluoromethanesulfonic Acid, *Anal. Biochem.*, 118:131 (1981).
- Evans et al., Design of Nonpeptidic Ligands for a Peptide Receptor: Cholecystokinin Antagonists, *J. Med. Chem.*, 30:1229 (1987).
- Fanslow et al., Structural characteristics of CD40 ligand that determine biological function, *Semin. Immunol.*, 6:267-278 (1994).
- Fauchere, Elements for the Rational Design of Peptide Drugs, *Adv. Drug Res.*, 15:29 (1986).
- Fieschi and Casanova, Mini Review: The role of interleukin-12 in human infectious diseases: only a faint signature, *Eur. J. Immunol.*, 33:1461-4 (2003).
- Fishwild et al., High-avidity human IgGk monoclonal antibodies from a novel strain of minilocus transgenic mice, *Nature Biotechnology*, 14:845-85 (1996).
- Fredericks et al., Identification of potent human anti-IL-1RI antagonist antibodies, *Protein Engineering, Design & Selection*, 17:95-106 (2004).
- GenBank Accession No. AB030000.
- GenBank Accession No. M65272.
- GenBank Accession No. NM 005535.
- GenBank Accession No. NM 144701.
- Ghilardi et al., Compromised Humoral and Delayed-Type Hypersensitivity Responses in IL-23-Deficient Mice, *The Journal of Immunology*, 172:2827-2833 (2004).
- Gribskov et al., Profile Analysis, *Meth. Enzym.*, 183:146-159 (1990).
- Gribskov et al., Profile analysis: Detection of distantly related proteins, *Proc. Nat. Acad. Sci.*, 84:4355-4358 (1987).
- Grosschedl et al., Introduction of a μ Immunoglobulin Gene into the Mouse Germ Line: Specific Expression in Lymphoid Cells and Synthesis of Functional Antibody, *Cell*, 38:647-658 (1984).
- Guisti et al., Somatic diversification of S107 from an antiphosphocholine to an anti-DNA autoantibody is due to a single base change in its heavy chain variable region, *Proc. Natl. Acad. Sci. USA*, 84(9):2926-30 (1987).
- Hammer et al., Diversity of alpha-fetoprotein gene expression in mice is generated by a combination of separate enhancer elements, *Science*, 253:53-58 (1987).
- Hanahan, Heritable formation of pancreatic β -cell tumours in transgenic mice expressing recombinant insulin/simian virus 40 oncogenes, *Nature*, 315:115-122 (1985).
- Happel et al., Divergent roles of IL-23 and IL-12 in host defense against *Klebsiella pneumoniae*, *The Journal of Experimental Medicine*, 202:761-769 (2005).
- Harding and Lonberg, Class Switching in Human Immunoglobulin Transgenic Mice, *Ann. N. Y. Acad. Sci.*, 764:536-546 (1995).
- Hayden et al., Single-chain mono- and bispecific antibody derivatives with novel biological properties and antitumour activity from a COS cell transient expression system, *Therapeutic Immunology*, 1:3-15 (1994).
- Held et al., Generation of a Protective T-Cell Response Following Coronavirus Infection of the Central Nervous System Is Not Dependent on IL-12/23 Signaling, *Viral Immunology*, 21:173-187 (2008).
- Henikoff et al., Amino acid substitution matrices from protein blocks, *Proc. Natl. Acad. Sci. U.S.A.*, 89:10915-10919 (1992).
- Hillyer et al., Investigating the role of the interleukin-23/-17A axis in rheumatoid arthritis, *Rheumatology*, 48:1581-1589 (2009).
- Hoeve et al., Divergent effects of IL-12 and IL-23 on the production of IL-17 by human T cells, *Eur. J. Immunol.*, 36:1-10 (2006).
- Holliger et al., Diabodies: Small bivalent and bispecific antibody fragments, *Proc. Natl. Acad. Sci. USA*, 90:6444-48 (1993).
- Holm et al., Protein folds and families: sequence and structure alignments, *Nucl. Acid. Res.*, 27:244-247 (1999).
- Holt et al., Domain antibodies: proteins for therapy, *Trends Biotechnol.*, 21(11):484-90 (2003).

(56)

References Cited

OTHER PUBLICATIONS

- Holtta et al., IL-23/IL-17 Immunity as a Hallmark of Crohn's Disease, *Inflamm Bowel Dis.*, 14:1175-1184 (2008).
- Honegger and Pluckthun, Yet another numbering scheme for immunoglobulin variable domains: an automatic modeling and analysis tool, *J. Mol. Biol.*, 309(3):657-670 (2001).
- Hoogenboom et al., By-passing Immunisation Human Antibodies from Synthetic Repertoires of Germline VH Gene Segments Rearranged in Vitro, *J. Mol. Biol.*, 227:381 (1991).
- Hopp et al., A Short Polypeptide Marker Sequence Useful for Recombinant Protein Identification and Purification, *Bio/Technology*, 6:1204 (1988).
- Hoppe et al., A parallel three stranded α -helical bundle at the nucleation site of collagen triple-helix formation, *FEBS Letters*, 344:191 (1994).
- Huston et al., Protein engineering of antibody binding sites: recovery of specific activity in an anti-digoxin single-chain Fv analogue produced in *Escherichia coli*, *Proc. Natl. Acad. Sci. U.S.A.*, 85:5879 (1988).
- Jakobovits et al., Analysis of homozygous mutant chimeric mice: Deletion of the immunoglobulin heavy-chain joining region blocks B-cell development and antibody production, *Proc. Natl. Acad. Sci. USA*, 90:2551-2555 (1993).
- Jakobovits et al., Germ-line transmission and expression of a human-derived yeast artificial chromosome, *Nature*, 362:255-258 (1993).
- Jones et al., Replacing the complementarity-determining regions in a human antibody with those from a mouse, *Nature*, 321 :522-525 (1986).
- Jones, Progress in protein structure prediction, *Curr. Opin. Struct. Biol.*, 7:377-387 (1997).
- Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication No. 91-3242 (1991).
- Kastelein, et al., Discovery and Biology of IL-23 and IL-27: Related but Functionally Distinct Regulators of Inflammation, *Annual Review of Immunology*, 25:221-42 (2007).
- Kelsey et al., Species- and tissue-specific expression of human α 1-antitrypsin in transgenic mice, *Genes and Devel.*, 1 :161-171 (1987).
- Khader et al., IL-23 Compensates for the Absence of IL-12p70 and Is Essential for the IL-17 Response during Tuberculosis but Is Dispensable for Protection and Antigen-Specific IFN- γ Responses if IL-12p70 Is Available, *The Journal of Immunology*, 175:788-795 (2005).
- Kirkland et al., Analysis of the fine specificity and cross-reactivity of monoclonal anti-lipid A antibodies, *J. Immunol.*, 137:3614-3619 (1986).
- Kollias et al., Regulated Expression of Human $\alpha\gamma$ -, β -, and Hybrid $\gamma\beta$ -Globin Genes in Transgenic Mice: Manipulation of the Developmental Expression Patterns, *Cell*, 46:89-94 (1986).
- Korndorfer et al., Crystallographic Analysis of an Anticalin With Tailored Specificity for fluorescein Reveals High Structural Plasticity of the Lipocalin Loop Region, *Proteins: Structure, Function, and Bioinformatics*, 53(1):121-9 (2003).
- Kortt et al., Dimeric and trimeric antibodies: high avidity scFvs for cancer targeting, *Biomol. Eng.*, 18:95-108 (2001).
- Kortt et al., Single-chain Fv fragments of anti-neuraminidase antibody NC10 containing five- and ten-residue linkers form dimers and with zero-residue linker a trimer, *Prot. Eng.*, 10:423 (1997).
- Kostelny et al., Formation of a bispecific antibody by the use of leucine zippers, *J. Immunol.*, 148:1547-1553 (1992).
- Kriangkum et al., Bispecific and bifunctional single chain recombinant antibodies, *Biomol. Eng.*, 18:31-40 (2001).
- Krumlauf et al., Developmental regulation of α -fetoprotein genes in transgenic mice, *Mol. Cell. Biol.*, 5:1639-1648 (1985).
- Kurzeja et al., New Interleukin-23 pathway inhibitors in dermatology, *Am. J. Clin. Dermatol.*, 12:113-125 (2011).
- Kussie et al., A single engineered amino acid substitution changes antibody fine specificity, *J. Immunol.*, 152(1):146-52 (1994).
- Kyte et al., A simple method for displaying the hydropathic character of a protein, *J. Mol. Biol.*, 157:105-131 (1982).
- Langowski et al., IL-23 promotes tumour incidence and growth, *Nature*, 442: 461-5 (2006).
- Lantto et al., Chain Shuffling to modify properties of recombinant immunoglobulins, *Methods Mol. Biol.*, 178:303-316 (2002).
- Leder et al., Consequences of widespread deregulation of the c-myc gene in transgenic mice: multiple neoplasms and normal development, *Cell*, 45:485-495 (1986).
- Lederman et al., A single amino acid substitution in a common African allele of the CD4 molecule ablates binding of the monoclonal antibody, OKT4, *Mol. Immunol.*, 28(11):1171-81 (1991).
- Lefranc et al., IMGT unique numbering for immunoglobulin and T cell receptor constant domain and Ig superfamily C-like domains, *Dev. Comp. Immunol.*, 29:185-203 (2005).
- Li et al., Temporal associations between interleukin 22 and the extracellular domains of IL-22R and IL-10R2, *Int. Immunopharmacol.*, 4(5):693-708 (2004).
- Liu et al., Fine mapping of the antigen-antibody interaction of scFv215, a recombinant antibody inhibiting RNA polymerase II from *Drosophila melanogaster*, *J. Mol. Recognit.*, 12(2):103-11 (1999).
- Liu, et al., Interleukin (IL)-23 p19 expression induced by IL-1 β in human fibroblast-like synoviocytes with rheumatoid arthritis via active nuclear factor-kB and AP-1 dependent pathway, *Rheumatology*, 46(8):1266-73 (2007).
- Lonberg and Huszar, Human Antibodies from Transgenic Mice, *Intern. Rev. Immunol.*, 13: 65-93 (1995).
- Lonberg et al., Antigen-specific human antibodies from mice comprising four distinct genetic modifications, *Nature*, 368:856-859 (1994).
- Lonberg et al., Fully human antibodies from transgenic mouse and phage display platforms, *Current Opinion in Immunology*, 20(4):450-459 (2008).
- Lonberg, Chapter 3, Transgenic Approaches to Human Monoclonal Antibodies, *Handbook of Exp. Pharmacology*, 113:49-101 (1994).
- Lupardus and Garcia, The Structure of Interleukin-23 Reveals the Molecular Basis of p40 subunit sharing with Interleukin-12, *J. Mol. Biol.*, 382:931-941 (2008).
- Mabry et al., Engineering of stable bispecific antibodies targeting IL-17A and IL23, *Protein Engineering Design & Selection*, 23(3):115-127 (2010).
- MacDonald, Expression of the pancreatic elastase I gene in transgenic mice, *Hepatology*, 7:425-515 (1987).
- Magram et al., Developmental regulation of a cloned adult β -globin gene in transgenic mice, *Nature*, 315:338-340 (1985).
- Marks et al., By-Passing Immunization: building high affinity human antibodies by chain shuffling, *BioTechnology*, 10:779 (1992).
- Marks et al., Human Antibodies from V-gene Libraries Displayed on Phage, *J. Mol. Biol.*, 222:581 (1991).
- Mason et al., The hypogonadal mouse: reproductive functions restored by gene therapy, *Science*, 234:1372-1378 (1986).
- Maynard et al., Antibody engineering, *Annu. Rev. Biomed. Eng.*, 2:339-76 (2000).
- Meeran et al., Interleukin-12-deficient mice are at greater risk of UV radiation-induced skin tumors and malignant transformation of papillomas to carcinomas, *Mol. Cancer Ther.*, 5: 825-32 (2006).
- Mendez et al., Functional transplant of megabase human immunoglobulin loci recapitulates human antibody response in mice, *Nature Genetics*, 15:146-156 (1997).
- Moldenhauer et al., Identity of HML-1 antigen on intestinal intraepithelial T cells and of B-ly7 antigen on hairy cell leukaemia, *Scand. J. Immunol.*, 32:77-82 (1990).
- Morel et al., Monoclonal antibodies to bovine serum albumin: affinity and specificity determinations, *Molec. Immunol.*, 25:7-15 (1988).
- Morrison et al., Chimeric human antibody molecules: mouse antigen-binding domains with human constant region domains, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1985).
- Moult, The current state of the art in protein structure prediction, *Curr. Op. in Biotech.*, 7:422-427 (1996).

(56)

References Cited

OTHER PUBLICATIONS

- Needleman et al., A general method applicable to the search for similarities in the amino acid sequence of two proteins, *J. Mol. Biol.*, 48:443-453 (1970).
- Oppmann et al., Novel p19 protein engages IL-12p40 to Form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12, *Immunity*, 13:715-725 (2000).
- Ornitz et al., Elastase I Promoter Directs Expression of Human Growth Hormone and SV40 T Antigen Genes to Pancreatic Acinar Cells in Transgenic Mice, Cold Spring Harbor Symp. *Quant. Biol.*, 50:399-409 (1986).
- Panka et al., Variable region framework differences result in decreased or increased affinity of variant anti-digoxin antibodies, *Proc. Natl. Acad. Sci. USA*, 85(9):3080-4 (1988).
- Parham et al., A receptor for the heterodimeric cytokine IL-23 is composed of IL-12R β 1 and a novel cytokine receptor subunit, IL-23R1, *J. Immunol.*, 168:5699-708 (2002).
- Pini et al., Design and use of a phage display library. Human antibodies with subnanomolar affinity against a marker of angiogenesis eluted from a two-dimensional gel, *J. Biol. Chem.*, 273(34):21769-76 (1998).
- Pinkert et al., An albumin enhancer located 10 kb upstream functions along with its promoter to direct efficient, liver-specific expression in transgenic mice, *Genes and Devel.*, 1:268-276 (1987).
- Poljak et al., Production and structure of diabodies, *Structure*, 2:1121-23 (1994).
- Powers et al., Expression of single-chain Fv-Fc fusions in *Pichia pastoris*, *Journal of Immunological Methods*, 251:123-135 (2001).
- Readhead et al., Expression of a myelin basic protein gene in transgenic shiverer mice: correction of the dysmyelinating phenotype, *Cell*, 48:703-712 (1987).
- Riechmann et al., Reshaping human antibodies for therapy, *Nature*, 332:323-27 (1988).
- Rizo and Gierasch, Constrained peptides: models of bioactive peptides and protein substructures, *Ann. Rev. Biochem.*, 61:387 (1992).
- Roque et al., Antibodies and genetically engineered related molecules: production and purification, *Biotechnol. Prog.*, 20:639-654 (2004).
- Rudikoff et al., Single amino acid substitution altering antigen-binding specificity, *Proc. Natl. Acad. Sci. USA*, 79(6):1979-83 (1982).
- Sani, Tissue-specific expression of rat myosin light-chain 2 gene in transgenic mice, *Nature*, 314:283-286 (1985).
- Schildbach et al., Contribution of a single heavy chain residue to specificity of an anti-digoxin monoclonal antibody, *Protein Sci.*, 3(5):737-49 (1994).
- Schildbach et al., Heavy chain position 50 is a determinant of affinity and specificity for the anti-digoxin antibody 26-10, *J. Biol. Chem.*, 268(29):21739-47 (1993).
- Shu et al., Secretion of a single-gene-encoded immunoglobulin from myeloma cells, *Proc. Natl. Acad. Sci. USA*, 90:7995-7999 (1993).
- Sippl et al., Threading thrills and threats, *Structure*, 4:15-19 (1996).
- Sojar et al., A chemical method for the deglycosylation of proteins, *Arch. Biochem. Biophys.*, 259:52 (1987).
- Songsivilai and Lachmann, Bispecific antibody: a tool for diagnosis and treatment of disease, *Clin. Exp. Immunol.*, 79:315-321 (1990).
- Stahli et al., Distinction of Epitopes by Monoclonal Antibodies, *Methods in Enzymology*, 92:242-253 (1983).
- Swift et al., Tissue-Specific Expression of the Rat Pancreatic Elastase / Gene in Transgenic Mice, *Cell*, 38:639-646 (1984).
- Taylor et al., A transgenic mouse that expresses a diversity of human sequence heavy and light chain immunoglobulins, *Nucleic Acids Research*, 20:6287-6295 (1992).
- Taylor et al., Human immunoglobulin transgenes undergo rearrangement, somatic mutation and class switching in mice that lack endogenous IgM, *International Immunology*, 6:579-591 (1994).
- Thomsen et al., Promoter-regulatory region of the major immediate early gene of human cytomegalovirus, *Proc. Natl. Acad. U.S.A.*, 81:659-663 (1984).
- Thornton et al., Prediction of progress at last, *Nature*, 354:105 (1991).
- Thotakura et al., Enzymatic deglycosylation of glycoproteins, *Meth. Enzymol.*, 138:350 (1987).
- Tonel et al., Cutting edge: A critical functional role for IL-23 in psoriasis, *J. Immunol.*, 185(10):5688-91 (2010).
- Tuaillon et al., Biased utilization of DHQ52 and JH4 Gene Segments in a Human Ig Transgenic Minilocus is Independent of Antigenic Selection, *J. Immunol.*, 152:2912-2920 (1994).
- Vajdos et al., Comprehensive functional maps of the antigen-binding site of an anti-ErbB2 antibody obtained with shotgun scanning mutagenesis, *J. Mol. Biol.*, 230(2):415-28 (2002).
- Veber and Freidinger, The design of metabolically-stable peptide analogs, *TINS*, p. 392-396 (1985).
- Verhoeven et al., Reshaping human antibodies: grafting an antilysozyme activity, *Science*, 239:1534-1536 (1988).
- Villa-Kamaroff et al., A bacterial clone synthesizing proinsulin, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-3731 (1978).
- Wagner et al., Nucleotide sequence of the thymidine kinase gene of herpes simplex virus type 1, *Proc. Natl. Acad. Sci. U.S.A.*, 78:1444-1445 (1981).
- Ward et al., Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*, *Nature*, 334:544 (1989).
- Xiang et al., Study of B72.3 combining sites by molecular modeling and site-directed mutagenesis, *Protein Eng.*, 13(5):339-44 (2000).
- Yago et al., IL-23 induces human osteoclastogenesis via IL-17 in vitro, and anti-IL-23 antibody attenuates collagen-induced arthritis in rats, *Arthritis Res and Ther.*, 9(5):R96 (2007).
- Yamamoto et al., Identification of a functional promoter in the long terminal repeat of rous sarcoma virus, *Cell*, 22:787-797 (1980).
- Cargill et al., A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes, *Am. J. Hum. Genet.*, 80(2):273-90 (2007).
- Bandzar et al., Crohn's disease: a review of treatment options and current research, *Cell Immunol.*, 286(1-2):45-52 (2013).
- Duvallet et al., Interleukin-23: A key cytokine in inflammatory diseases, *Ann. Med.*, 43(7):503-11 (2011).
- Fitch et al., Pathophysiology of psoriasis: recent advances on IL-23 and Th17 cytokines, *Curr Rheumatol Rep.*, 9(6):461-4 (2007).
- Griffiths et al., Pathogenesis and clinical features of psoriasis, *Lancet*, 370(9583):263-71 (2007).
- Harlow et al., Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990).
- Hue et al., Interleukin-23 drives innate and T cell-mediated intestinal inflammation, *J. Exp. Med.*, 203(11):2473-83 (2006).
- Kullberg et al., IL-23 plays a key role in *Helicobacter hepaticus*-induced T cell-dependent colitis, *J. Exp. Med.*, 203(11):2485-94 (2006).
- Lee et al., Increased expression of interleukin 23 p19 and p40 in lesional skin of patients with psoriasis vulgaris, *J. Exp. Med.*, 199(1):125-30 (2004).
- Leonardi et al., Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 1), *Lancet*, 371(9625):1665-74 (2008).
- Lowes et al., Pathogenesis and therapy of psoriasis, *Nature*, 445(7130):866-73 (2007).
- Nestle et al., Psoriasis, *N. Engl. J. Med.*, 361(5):496-509 (2009).
- Neurath, IL-23: a master regulator in Crohn disease, *Nat. Med.*, 13(1):26-8 (2007).
- Nickoloff et al., Recent insights into the immunopathogenesis of psoriasis provide new therapeutic opportunities, *J. Clin. Invest.*, 113(12):1664-75 (2004).
- Papp et al., Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 52-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 2), *Lancet*, 371(9625):1675-84 (2008).
- Perrier et al., Cytokine blockade in inflammatory bowel diseases, *Immunotherapy*, 3(11):1341-52 (2011).

(56)

References Cited

OTHER PUBLICATIONS

Piskin et al., In vitro and in situ expression of IL-23 by keratinocytes in healthy skin and psoriasis lesions: enhanced expression in psoriatic skin, *J. Immunol.*, 176(3):1908-15 (2006).

Torti et al., Interleukin-12, interleukin-23, and psoriasis: Current prospects, *J. Amer. Acad. Derm.*, 57(6):1059-68 (2007).




Fig. 1A




Fig. 1B

1

HUMAN IL-23 ANTIGEN BINDING PROTEINS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a divisional application of U.S. patent application Ser. No. 14/228,556 filed Mar. 28, 2014 (now U.S. Pat. No. 9,487,580) which is a continuation of U.S. patent application Ser. No. 13/504,449 filed Aug. 31, 2012 (now U.S. Pat. No. 8,722,033), which in turn is a United States National Stage Application of PCT/US10/54148, filed Oct. 26, 2010, which in turn claims the benefit under 35 U.S.C. 119(e) of U.S. Provisional Application No. 61/254,982, filed Oct. 26, 2009 and U.S. Provisional Application No. 61/381,287, filed Sep. 9, 2010, which are incorporated herein by reference.

BACKGROUND

Interleukin 23 (IL-23), a heterodimeric cytokine, is a potent inducer of pro-inflammatory cytokines. IL-23 is related to the heterodimeric cytokine Interleukin 12 (IL-12) both sharing a common p40 subunit. In IL-23, a unique p19 subunit is covalently bound to the p40 subunit. In IL-12, the unique subunit is p35 (Oppmann et al., *Immunity*, 2000, 13: 713-715). The IL-23 heterodimeric protein is secreted. Like IL-12, IL-23 is expressed by antigen presenting cells (such as dendritic cells and macrophages) in response to activation stimuli such as CD40 ligation, Toll-like receptor agonists and pathogens. IL-23 binds a heterodimeric receptor comprising an IL-12R β 1 subunit (which is shared with the IL-12 receptor) and a unique receptor subunit, IL-23R. The IL-12 receptor consists of IL-12R β 1 and IL-12R β 2. IL-23 binds its heterodimeric receptor and signals through JAK2 and Tyk2 to activate STAT1, 3, 4 and 5 (Parham et al., *J. Immunol.* 2002, 168:5699-708). The subunits of the receptor are predominantly co-expressed on activated or memory T cells and natural killer cells and also at lower levels on dendritic cells, monocytes, macrophages, microglia, keratinocytes and synovial fibroblasts. IL-23 and IL-12 act on different T cell subsets and play substantially different roles in vivo.

IL-23 acts on activated and memory T cells and promotes survival and expansion of the T cell subset, Th17. Th17 cells produce proinflammatory cytokines including IL-6, IL-17, TNF α , IL-22 and GM-CSF. IL-23 also acts on natural killer cells, dendritic cells and macrophages to induce pro-inflammatory cytokine expression. Unlike IL-23, IL-12 induces the differentiation of naïve CD4⁺ T cells into mature Th1 IFN γ -producing effector cells, and induces NK and cytotoxic T cell function by stimulating IFN γ production. Th1 cells driven by IL-12 were previously thought to be the pathogenic T cell subset in many autoimmune diseases, however, more recent animal studies in models of inflammatory bowel disease, psoriasis, inflammatory arthritis and multiple sclerosis, in which the individual contributions of IL-12 versus IL-23 were evaluated have firmly established that IL-23, not IL-12, is the key driver in autoimmune/inflammatory disease (Ahern et al., *Immun. Rev.* 2008 226:147-159; Cua et al., *Nature* 2003 421:744-748; Yago et al., *Arthritis Res and Ther.* 2007 9(5): R96). It is believed that IL-12 plays a critical role in the development of protective innate and adaptive immune responses to many intracellular pathogens and viruses and in tumor immune surveillance. See Kastelein, et al., *Annual Review of Immunology*, 2007, 25: 221-42; Liu, et al., *Rheumatology*, 2007, 46(8): 1266-73; Bowman et al., *Current Opinion in Infec-*

2

tious Diseases, 2006 19:245-52; Fieschi and Casanova, *Eur. J. Immunol.* 2003 33:1461-4; Meeran et al., *Mol. Cancer Ther.* 2006 5: 825-32; Langowski et al., *Nature* 2006 442: 461-5. As such, IL-23 specific inhibition (sparing IL-12 or the shared p40 subunit) should have a potentially superior safety profile compared to dual inhibition of IL-12 and IL-23.

Therefore, use of IL-23 specific antagonists that inhibit human IL-23 (such as antibodies that bind at least the unique p19 subunit or bind both the p19 and p40 subunits of IL-23) that spare IL-12 should provide efficacy equal to or greater than IL-12 antagonists or p40 antagonists without the potential risks associated with inhibition of IL-12. Murine, humanized and phage display antibodies selected for inhibition of recombinant IL-23 have been described; see for example U.S. Pat. No. 7,491,391, WIPO Publications WO1999/05280, WO2007/0244846, WO2007/027714, WO 2007/076524, WO2007/147019, WO2008/103473, WO 2008/103432, WO2009/043933 and WO2009/082624. However, there is a need for fully human therapeutic agents that are able to inhibit native human IL-23. Such therapeutics are highly specific for the target, particularly in vivo. Complete inhibition of the in vivo target can result in lower dose formulations, less frequent and/or more effective dosing which in turn results in reduced cost and increased efficiency. The present invention provides such IL-23 antagonists.

SUMMARY

Antigen binding proteins that bind IL-23, particularly native human IL-23, are provided. The human IL-23 antigen binding proteins can reduce, inhibit, interfere with, and/or modulate at least one of the biological responses related to IL-23, and as such, are useful for ameliorating the effects of IL-23 related diseases or disorders. IL-23 antigen binding proteins can be used, for example, to reduce, inhibit, interfere with and/or modulate IL-23 signaling, IL-23 activation of Th17 cells, IL-23 activation of NK cells, or inducing production of proinflammatory cytokines.

Also provided are expression systems, including cell lines, for the production of IL-23 antigen binding proteins and methods of diagnosing and treating diseases related to human IL-23.

Some of the antigen binding proteins that bind IL-23 that are provided comprise at least one heavy chain variable region comprising a CDRH1, a CDRH2 and a CDRH3 selected from the group consisting of: a CDRH1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH1 as shown in TABLE 3; a CDRH2 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRH2 as shown in TABLE 3; a CDRH3 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRH3 as shown in TABLE 3; and comprising at least one light chain variable region comprising a CDRL1, a CDRL2 and a CDRL3 selected from the group consisting of: a CDRL1 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRL1 as shown in TABLE 3; a CDRL2 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL2 as shown in TABLE 3; a CDRL3 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL3 as shown in TABLE 3. In one embodiment is provided isolated antigen binding proteins comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 94, 97, 100, and 103;

a CDRH2 selected from the group consisting of SEQ ID NO:92, 95, 98, 101, 104, 107, and 110; a CDRH3 selected from the group consisting of SEQ ID NO: 93, 96, 99, 102, and 105; a CDRL1 selected from the group consisting of SEQ ID NO: 62, 65, 68, 71, and 74; a CDRL2 selected from the group consisting of SEQ ID NO:63, 66, 69, 72, 75, and 78; and a CDRL3 selected from the group consisting of SEQ ID NO:64, 67, 70 and 73. In another embodiment is provided isolated antigen bindings protein of comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 106, 109, 112, and 115; a CDRH2 selected from the group consisting of SEQ ID NO: 113, 116, 118, 120, 121, and 122; a CDRH3 selected from the group consisting of SEQ ID NO: 108, 111, 114, 117, and 119; a CDRL1 selected from the group consisting of SEQ ID NO: 77, 80, 83, 85, 86, 87, 88, 89 and 90; a CDRL2 is SEQ ID NO: 81; and a CDRL3 selected from the group consisting of SEQ ID NO: 76, 79, 82 and 84. In another embodiment is provided an isolated antigen-binding protein of that comprises at least one heavy chain variable region and at least one light chain variable region. In yet another embodiment is provided an isolated antigen-binding protein as described above that comprise at least two heavy chain variable regions and at least two light chain variable regions. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is coupled to a labeling group.

Also provided are isolated antigen binding proteins that bind IL-23 selected from the group consisting of a) an antigen binding protein having CDRH1 of SEQ ID NO:129, CDRH2 of SEQ ID NO:132, CDRH3 of SEQ ID NO:136, and CDRL1 of SEQ ID NO:123, CDRL2 of SEQ ID NO:81, and CDRL3 of SEQ ID NO: 76; b) an antigen binding protein having CDRH1 of SEQ ID NO:131, CDRH2 of SEQ ID NO: 134, CDRH3 of SEQ ID NO:137 and CDRL1 of SEQ ID NO:124, CDRL2 of SEQ ID NO:126 and CDRL3 of SEQ ID NO:128; c) a) an antigen binding protein having CDRH1 of SEQ ID NO:130, CDRH2 of SEQ ID NO:133, CDRH3 of SEQ ID NO:99 and CDRL1 of SEQ ID NO:68, CDRL2 of SEQ ID NO:69, and CDRL3 of SEQ ID NO:67; and d) an antigen binding protein having CDRH1 SEQ ID NO:91, CDRH2 SEQ ID NO: 135, CDRH3 SEQ ID NO:138 and CDRL1 SEQ ID NO:125, CDRL2 SEQ ID NO:127, and CDRL3 SEQ ID NO:64.

Also provided are isolated antigen binding proteins that bind IL-23 comprising at least one heavy chain variable region and at least one light chain variable region, selected from the group consisting of: a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-113 of SEQ ID NO:31; and a light chain variable region comprising amino acid residues 23-36, 52-58 and 91-101 of SEQ ID NO:1; a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-110 of SEQ ID NO:34 and heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-110 of SEQ ID NO:36; and a light chain variable region comprising amino acid residues 23-36, 52-62 and 97-105 of SEQ ID NO:4; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:38; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:7; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:40; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 94-106 of SEQ ID NO:9; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:42; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:11; a heavy chain variable

region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO:44; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:13; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:46 or SEQ ID NO:153; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:15; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:48; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:17; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 101-109 of SEQ ID NO:50; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:19; a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO: 52; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 98-107 of SEQ ID NO:21; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:54; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:23; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:56; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:25; and a heavy chain variable region comprising amino acid residues 31-37, 52-57 and 100-109 of SEQ ID NO:58; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:27.

Provided herein is an isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region sequence differs by no more than 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions, additions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2; and wherein the light chain variable region sequence differs by no more than 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions, additions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

Also provided is an isolated antigen binding protein that binds IL-23 selected from the group consisting of a) a heavy chain variable region of SEQ ID NO:140 and a light chain variable region of SEQ ID NO: 30; b) a heavy chain variable region of SEQ ID NO:141 and a light chain variable region of SEQ ID NO:61; c) a heavy chain variable region of SEQ ID NO:142 and a light chain variable region of SEQ ID NO:4; and d) a heavy chain variable region of SEQ ID NO:143 and a light chain variable region of SEQ ID NO:139.

Also provided is an isolated antigen binding protein comprising a heavy chain variable region comprising of an amino acid sequence having at least 90%, 95%, 96%, 97%, 98% or 99% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 90% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27. In another embodiment is an isolated antigen binding protein comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 44, 46, 48, 50, 52, 54, 56, 58 and 153, and a light chain variable region selected from the group consisting of SEQ ID NO:13, 15, 17, 19, 21, 23, 25, and 27. In yet another embodiment is an isolated antigen binding protein comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 31, 34, 36, 38, 40 and 42, and a

5

light chain variable region selected from the group consisting of SEQ ID NO: 1, 4, 7, 9 and 11.

Also provided is an isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region selected from the group consisting of: a) a heavy chain variable region of SEQ ID NO:31 and a light chain variable region of SEQ ID NO:1; b) a heavy chain variable region of SEQ ID NO:34 or 36 and a light chain variable region of SEQ ID NO:4; c) a heavy chain variable region of SEQ ID NO:38 and a light chain variable region of SEQ ID NO: 7; d) a heavy chain variable region of SEQ ID NO:40 and a light chain variable region of SEQ ID NO:9; e) a heavy chain variable region of SEQ ID NO:42 and a light chain variable region of SEQ ID NO: 11; f) a heavy chain variable region of SEQ ID NO:44 and a light chain variable region of SEQ ID NO:13; g) a heavy chain variable region of SEQ ID NO:46 or SEQ ID NO:153 and a light chain variable region of SEQ ID NO:15; h) a heavy chain variable region of SEQ ID NO:48 and a light chain variable region of SEQ ID NO:17; i) a heavy chain variable region of SEQ ID NO:50 and a light chain variable region of SEQ ID NO: 19; j) a heavy chain variable region of SEQ ID NO:52 and a light chain variable region of SEQ ID NO:21; k) a heavy chain variable region of SEQ ID NO:54 and a light chain variable region of SEQ ID NO:23; l) a heavy chain variable region of SEQ ID NO:56 and a light chain variable region of SEQ ID NO:25; and m) a heavy chain variable region of SEQ ID NO:58 and a light chain variable region of SEQ ID NO:27.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 30, 31, 32, 49, 50, 52, 53, 56, 92 and 94 of SEQ ID NO:15, wherein the residue contacts have a difference value of greater than or equal to $10 \text{ \AA}^2$ as determined by solvent exposed surface area. Within one embodiment the residue contacts comprise residues 31-35, 54, 58-60, 66, and 101-105 of SEQ ID NO:46.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 31-34, 51, 52, 55, 68, 93 and 98 of SEQ ID NO:1, wherein the residue contacts have a difference value of greater than or equal to $10 \text{ \AA}^2$ as determined by solvent exposed surface area. Within one embodiment the residue contacts comprise residues 1, 26, 28, 31, 32, 52, 53, 59, 76, 101, 102 and 104-108 of SEQ ID NO:31.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 32-35, 54, 58-60, 66 and 101-105 of SEQ ID NO:46, as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 31-35, 54, 56, 58-60, 66 and 101-105 of SEQ ID NO:46.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 30-32, 49, 52, 53, 91-94 and 96 of SEQ ID NO:15, as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 30-32, 49, 50, 52, 53, 56, 91-94 and 96 of SEQ ID NO:15.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 26-28, 31, 53, 59, 102 and

6

104-108 of SEQ ID NO:31, as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 1, 26-28, 30-32, 52, 53, 59, 100, and 102-108 of SEQ ID NO:31.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 Å or less from residues 31-34, 51, 52, 55, 68 and 93 of SEQ ID NO:1 as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 29, 31-34, 51, 52, 55, 68, 93 and 100 of SEQ ID NO:1.

Also provided is an isolated antigen binding protein as described above, wherein the antigen binding protein is an antibody. In one embodiment is provided an isolated antigen binding protein wherein the antibody is a monoclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, a chimeric antibody, a multispecific antibody, or an antibody fragment thereof. In another embodiment is provided an isolated antigen binding protein wherein the antibody fragment is a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, a Fv fragment, a diabody, or a single chain antibody molecule. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is a human antibody. In still another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is a monoclonal antibody. In another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is of the IgG1-, IgG2-IgG3- or IgG4-type. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is of the IgG1- or IgG2-type.

An isolated nucleic acid molecule encoding an antigen binding protein as described above, is also provided. In one embodiment is provided an isolated nucleic acid molecule wherein at least one heavy chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:32, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59 and 152 and at least one light chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:2, 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, and 28. In another embodiment is provided a nucleic acid molecule wherein the nucleic acid molecule is operably linked to a control sequence. In another embodiment is provided a vector comprising a nucleic acid molecule as described above. In yet another embodiment is provided a host cell comprising the nucleic acid molecule as described above. In another embodiment is provided a host cell comprising the vector described above. In yet another embodiment is provided an isolated polynucleotide sufficient for use as a hybridization probe, PCR primer or sequencing primer that is a fragment of the nucleic acid molecule as described above or its complement.

Also provided is a method of making the antigen binding protein as described above, comprising the step of preparing said antigen binding protein from a host cell that secretes said antigen binding protein.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises a residue contact within residues 46-58, a residue contact within residues 112-120, and a residue contact within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, wherein the residue contact has a difference value greater than or equal to $10 \text{ \AA}^2$ as determined by solvent exposed surface area. In one embodi-

ment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or thirteen residue contacts within residues 46-58, one, two, three, four, five, six, seven, eight, nine or ten residue contacts within residues 112-120, and one, two, three, four, five, six, seven, eight or nine residue contacts within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145. In another embodiment is provided wherein the covered patch formed when the antigen binding protein binds to human IL-23 comprises a residue contact within residues 121-125 of the human IL-23p40 subunit as described in SEQ ID NO:147. In a related embodiment is wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises one, two, three, four or five residue contacts within residues 121-125 of the human IL-23p40 subunit as described in SEQ ID NO:147. Within another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 46, 47, 49, 50, 53, 112-116, 118, 120, 155, 156, 159, 160, and 163 of SEQ ID NO:145. In another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 46, 47, 49, 50, 53, 112-118, 120, 155, 156, 159, 160, and 163 of SEQ ID NO:145. Within another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residues 46, 47, 49, 50, 53-55, 57, 58, 112-116, 118-120, 155, 156, 159, 160, 162 and 163 of SEQ ID NO:145. In a related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 122 of the human IL-23p40 subunit as described in SEQ ID NO:147. In another related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 122 and 124 of the human IL-23p40 subunit as described in SEQ ID NO:147. In yet another related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 121-123 and 125 of the human IL-23p40 subunit as described in SEQ ID NO:147. In a further related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 121-123, 125 and 283 of the human IL-23p40 subunit as described in SEQ ID NO:147.

Also provided is an isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23 said antigen binding protein is 5 Å or less from a residue within residues 46-58, from a residue within residues 112-123, and from a residue within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, as determined by X-ray crystallography. In one embodiment, when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or thirteen residues within residues 46-58, from one, two, three, four, five, six, seven, eight, nine or ten, residues within residues 112-123, and from one, two, three, four, five, six, seven, eight or nine residues within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145. Within another embodiment when the antigen binding protein is bound to human IL-23 the antigen binding protein is 5 Å or less from residues 46-50, 113-116, 120, 156, 159, 160 and 163 of SEQ ID NO:145. Within another embodiment when the antigen

binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 112-120, 156, 159, 160 and 163 of SEQ ID NO:145. Within a related embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 53, 112-120, 156, 159, 160 and 163 of SEQ ID NO:145. Within another embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 53-55, 58, 113-116, 120, 121, 156, 159, 160, 162 and 163 of SEQ ID NO:145. Within a related embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-51, 53-55, 57, 58, 112-116, 118-121, 123, 155, 156, 159, 160, 162 and 163 of SEQ ID NO:145. Within a further embodiment when the antigen binding protein is bound to human IL-23 the antigen binding protein is 5 Å or less from a residue within residues 121-125, of the human IL-23p40 subunit as described in SEQ ID NO:147, as determined by X-ray crystallography. With a related embodiment when the antigen binding protein is bound to human IL-23, said antigen binding protein is 5 Å or less from residues 122 and 124 of SEQ ID NO:147. Within another embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 121-123 and 125 of SEQ ID NO:147.

Also provided is an isolated antigen binding protein as described above, wherein the antigen binding protein has at least one property selected from the group consisting of: a) reducing human IL-23 activity; b) reducing production of a proinflammatory cytokine; c) binding to human IL-23 with a KD of $\leq 5 \times 10^{-8}$ M; d) having a Koff rate of $\leq 5 \times 10^{-6}$ 1/s; and d) having an IC50 of ≤ 400 pM.

A pharmaceutical composition comprising at least one antigen binding protein as described above and pharmaceutically acceptable excipient is provided. In one embodiment is provided a pharmaceutical composition further comprises a labeling group or an effector group. In yet another embodiment is provided a pharmaceutical composition wherein the labeling group is selected from the group consisting of isotopic labels, magnetic labels, redox active moieties, optical dyes, biotinylated groups and predetermined polypeptide epitopes recognized by a secondary reporter. In yet another embodiment is provided a pharmaceutical composition wherein the effector group is selected from the group consisting of a radioisotope, radionuclide, a toxin, a therapeutic group and a chemotherapeutic group.

Also provided is a method for treating or preventing a condition associated with IL-23 in a patient, comprising administering to a patient in need thereof an effective amount of at least one isolated antigen binding protein as described above. In one embodiment is provided a method of wherein the condition is selected from the group consisting of an inflammatory disorder, a rheumatic disorder, an autoimmune disorder, an oncological disorder and a gastrointestinal disorder. In yet another embodiment is provided a method wherein the condition is selected from the group consisting of multiple sclerosis, rheumatoid arthritis, cancer, psoriasis, inflammatory bowel disease, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, psoriatic arthritis, autoimmune myocarditis; type 1 diabetes and ankylosing spondylitis. In still another embodiment is provided a method wherein the isolated antigen-binding protein is administered alone or as a combination therapy.

Also provided is a method of reducing IL-23 activity in a patient comprising administering an effective amount of at least one antigen binding protein as described above. In one

istry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques can be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

All patents and other publications identified are expressly incorporated herein by reference in their entirety for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with information described herein.

The polynucleotide and protein sequences of the p19 subunit of human IL-23 (SEQ ID NOS: 144 and 145), the shared p40 subunit (SEQ ID NOS: 146 and 147), the human IL-23 receptor heterodimeric subunits IL-12R β 1 (SEQ ID NOS: 150 and 151) and IL-23R (SEQ ID NOS: 148 and 149), are known in the art, see for example, GenBank Accession Nos. AB030000; M65272, NM_005535, NM_144701, as are those from other mammalian species. Recombinant IL-23 and IL-23 receptor proteins including single chain and Fc proteins as well as cells expressing the IL-23 receptor have been described or are available from commercial sources. (see for example, Oppmann et al., *Immunity*, 2000, 13: 713-715; R&D Systems, Minneapolis, Minn.; United States Biological, Swampscott, Mass.; WIPO Publication No. WO 2007/076524). Native human IL-23 can be obtained from human cells such as dendritic cells using methods known in the art including those described herein.

IL-23 is a heterodimeric cytokine comprised of a unique p19 subunit that is covalently bound to a shared p40 subunit. The p19 subunit comprises four α -helices, "A", "B", "C" and "D" in an up-up-down-down motif joined by three intra-helix loops between the A and B helices, between the B and C helices and between the C and D helices, see Oppmann et al., *Immunity*, 2000, 13: 713-715 and Beyer, et al., *J Mol Biol*, 2008, 382(4): 942-55. The A and D helices of 4 helical bundle cytokines are believed to be involved with receptor binding. The p40 subunit comprises three β -sheet sandwich domains, D1, D2 and D3 (Lupardus and Garcia, *J. Mol. Biol.*, 2008, 382:931-941.

The term "polynucleotide" includes both single-stranded and double-stranded nucleic acids and includes genomic DNA, RNA, mRNA, cDNA, or synthetic origin or some combination thereof which is not associated with sequences normally found in nature. Isolated polynucleotides comprising specified sequences may include, in addition to the specified sequences, coding sequences for up to ten or even up to twenty other proteins or portions thereof, or may include operably linked regulatory sequences that control expression of the coding region of the recited nucleic acid sequences, and/or may include vector sequences. The nucleotides comprising the polynucleotide can be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. The modifications include base modifications such as bromouridine and inosine derivatives, ribose modifications such as 2',3'-dideoxyribose, and internucleotide linkage modifications such as phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphorothioanilthioate, phosphoraniladate and phosphoramidate.

The term “oligonucleotide” means a polynucleotide comprising 100 or fewer nucleotides. In some embodiments, oligonucleotides are 10 to 60 bases in length. In other embodiments, oligonucleotides are 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 nucleotides in length. Oligonucleotides may be single stranded or double stranded, e.g., for use in the construction of a mutant gene. Oligonucleotides may be sense or antisense oligonucleotides. An oligonucleotide can

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001) and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane *Antibodies: A Laboratory Manual* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The terminology used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chem-

include a detectable label, such as a radiolabel, a fluorescent label, a hapten or an antigenic label, for detection assays. Oligonucleotides may be used, for example, as PCR primers, cloning primers or hybridization probes.

The terms "polypeptide" or "protein" means a macromolecule having the amino acid sequence of a native protein, that is, a protein produced by a naturally-occurring and non-recombinant cell; or it is produced by a genetically-engineered or recombinant cell, and comprise molecules having the amino acid sequence of the native protein, or molecules having one or more deletions from, insertions to, and/or substitutions of the amino acid residues of the native sequence. The term also includes amino acid polymers in which one or more amino acids are chemical analogs of a corresponding naturally-occurring amino acid and polymers. The terms "polypeptide" and "protein" encompass IL-23 antigen binding proteins (such as antibodies) and sequences that have one or more deletions from, additions to, and/or substitutions of the amino acid residues of the antigen binding protein sequence. The term "polypeptide fragment" refers to a polypeptide that has an amino-terminal deletion, a carboxyl-terminal deletion, and/or an internal deletion as compared with the full-length native protein. Such fragments may also contain modified amino acids as compared with the native protein. In certain embodiments, fragments are about five to 500 amino acids long. For example, fragments may be at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 50, 70, 100, 110, 150, 200, 250, 300, 350, 400, or 450 amino acids long. Useful polypeptide fragments include immunologically functional fragments of antibodies, including binding domains. In the case of an IL-23 antigen binding protein, such as an antibody, useful fragments include but are not limited to one or more CDR regions, a variable domain of a heavy or light chain, a portion of an antibody chain, a portion of a variable region including less than three CDRs, and the like.

"Amino acid" includes its normal meaning in the art. The twenty naturally-occurring amino acids and their abbreviations follow conventional usage. See, *Immunology-A Synthesis*, 2nd Edition, (E. S. Golub and D. R. Gren, eds.), Sinauer Associates: Sunderland, Mass. (1991). Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as [alpha]-, [alpha]-disubstituted amino acids, N-alkyl amino acids, and other unconventional amino acids may also be suitable components for polypeptides. Examples of unconventional amino acids include: 4-hydroxyproline, [gamma]-carboxyglutamate, [epsilon]-N,N,N-trimethyllysine, [epsilon]-N-acetyllysine, O-phosphoserine, N-acetylserine, N-formyl-methionine, 3-methylhistidine, 5-hydroxylysine, [sigma]-N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxyl-terminal direction, in accordance with standard usage and convention.

The term "isolated protein" refers to a protein, such as an antigen binding protein (an example of which could be an antibody), that is purified from proteins or polypeptides or other contaminants that would interfere with its therapeutic, diagnostic, prophylactic, research or other use. As used herein, "substantially pure" means that the described species of molecule is the predominant species present, that is, on a molar basis it is more abundant than any other individual species in the same mixture. In certain embodiments, a substantially pure molecule is a composition wherein the object species comprises at least 50% (on a molar basis) of

all macromolecular species present. In other embodiments, a substantially pure composition will comprise at least 80%, 85%, 90%, 95%, or 99% of all macromolecular species present in the composition. In certain embodiments, an essentially homogeneous substance has been purified to such a degree that contaminating species cannot be detected in the composition by conventional detection methods and thus the composition consists of a single detectable macromolecular species.

A "variant" of a polypeptide (e.g., an antigen binding protein such as an antibody) comprises an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into the amino acid sequence relative to another polypeptide sequence. Variants include fusion proteins. A "derivative" of a polypeptide is a polypeptide that has been chemically modified in some manner distinct from insertion, deletion, or substitution variants, e.g., via conjugation to another chemical moiety.

The terms "naturally occurring" or "native" as used throughout the specification in connection with biological materials such as polypeptides, nucleic acids, host cells, and the like, refers to materials which are found in nature, such as native human IL-23. In certain aspects, recombinant antigen binding proteins that bind native IL-23 are provided. In this context, a "recombinant protein" is a protein made using recombinant techniques, i.e., through the expression of a recombinant nucleic acid as described herein. Methods and techniques for the production of recombinant proteins are well known in the art.

The term "antibody" refers to an intact immunoglobulin of any isotype, or a fragment thereof that can compete with the intact antibody for specific binding to the target antigen, and includes, for instance, chimeric, humanized, fully human, and bispecific antibodies. An antibody as such is a species of an antigen binding protein. Unless otherwise indicated, the term "antibody" includes, in addition to antibodies comprising two full-length heavy chains and two full-length light chains, derivatives, variants, fragments, and muteins thereof, examples of which are described below. An intact antibody generally will comprise at least two full-length heavy chains and two full-length light chains, but in some instances may include fewer chains such as antibodies naturally occurring in camelids which may comprise only heavy chains. Antibodies may be derived solely from a single source, or may be "chimeric," that is, different portions of the antibody may be derived from two different antibodies as described further below. The antigen binding proteins, antibodies, or binding fragments may be produced in hybridomas, by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies.

The term "functional fragment" (or simply "fragment") of an antibody or immunoglobulin chain (heavy or light chain), as used herein, is an antigen binding protein comprising a portion (regardless of how that portion is obtained or synthesized) of an antibody that lacks at least some of the amino acids present in a full-length chain but which is capable of specifically binding to an antigen. Such fragments are biologically active in that they bind specifically to the target antigen and can compete with other antigen binding proteins, including intact antibodies, for specific binding to a given epitope. In one aspect, such a fragment will retain at least one CDR present in the full-length light or heavy chain, and in some embodiments will comprise a single heavy chain and/or light chain or portion thereof. These biologically active fragments may be produced by recombinant DNA techniques, or may be produced by enzymatic or

chemical cleavage of antigen binding proteins, including intact antibodies. Fragments include, but are not limited to, immunologically functional fragments such as Fab, Fab', F(ab')₂, Fv, domain antibodies and single-chain antibodies, and may be derived from any mammalian source, including but not limited to human, mouse, rat, camelid or rabbit. It is contemplated further that a functional portion of the antigen binding proteins disclosed herein, for example, one or more CDRs, could be covalently bound to a second protein or to a small molecule to create a therapeutic agent directed to a particular target in the body, possessing bifunctional therapeutic properties, or having a prolonged serum half-life.

The term "compete" when used in the context of antigen binding proteins (e.g., neutralizing antigen binding proteins or neutralizing antibodies) means competition between antigen binding proteins as determined by an assay in which the antigen binding protein (e.g., antibody or immunologically functional fragment thereof) under test prevents or inhibits specific binding of a reference antigen binding protein (e.g., a ligand, or a reference antibody) to a common antigen (e.g., an IL-23 protein or a fragment thereof). Numerous types of competitive binding assays can be used, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see, e.g., Stahl et al., 1983, *Methods in Enzymology* 92:242-253); solid phase direct biotin-avidin EIA (see, e.g., Kirkland et al., 1986, *J. Immunol.* 137:3614-3619) solid phase direct labeled assay, solid phase direct labeled sandwich assay (see, e.g., Harlow and Lane, 1988, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using 1-125 label (see, e.g., Morel et al., 1988, *Molec. Immunol.* 25:7-15); solid phase direct biotin-avidin EIA (see, e.g., Cheung et al., 1990, *Virology* 176:546-552); and direct labeled RIA (Moldenhauer et al., 1990, *Scand. J. Immunol.* 32:77-82). Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabelled test antigen binding protein and a labeled reference antigen binding protein.

Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test antigen binding protein. Usually the test antigen binding protein is present in excess. Antigen binding proteins identified by competition assay (competing antigen binding proteins) include antigen binding proteins binding to the same epitope as the reference antigen binding proteins and antigen binding proteins binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antigen binding protein for steric hindrance to occur. Usually, when a competing antigen binding protein is present in excess, it will inhibit specific binding of a reference antigen binding protein to a common antigen by at least 40%, 45%, 50%, 55%, 60%, 65%, 70% or 75%. In some instance, binding is inhibited by at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% 98%, 99% or more.

The term "epitope" or "antigenic determinant" refers to a site on an antigen to which an antigen binding protein binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. Epitope determinants may include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl groups, and may have specific three dimensional structural characteristics, and/or specific charge char-

acteristics. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35 amino acids in a unique spatial conformation. Epitopes can be determined using methods known in the art.

IL-23 Antigen Binding Proteins

An "antigen binding protein" as used herein means a protein that specifically binds a specified target antigen; the antigen as provided herein is IL-23, particularly human IL-23, including native human IL-23. Antigen binding proteins as provided herein interact with at least a portion of the unique p19 subunit of IL-23, detectably binding IL-23; but do not bind with any significance to IL-12 (e.g., the p40 and/or the p35 subunits of IL-12), thus "sparing IL-12". As a consequence, the antigen binding proteins provided herein are capable of impacting IL-23 activity without the potential risks that inhibition of IL-12 or the shared p40 subunit might incur. The antigen binding proteins may impact the ability of IL-23 to interact with its receptor, for example by impacting binding to the receptor, such as by interfering with receptor association. In particular, such antigen binding proteins totally or partially reduce, inhibit, interfere with or modulate one or more biological activities of IL-23. Such inhibition or neutralization disrupts a biological response in the presence of the antigen binding protein compared to the response in the absence of the antigen binding protein and can be determined using assays known in the art and described herein. Antigen binding proteins provided herein inhibit IL-23-induced proinflammatory cytokine production, for example IL-23-induced IL-22 production in whole blood cells and IL-23-induced IFN γ expression in NK and whole blood cells. Reduction of biological activity can be about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% 98%, 99% or more.

An antigen binding protein may comprise a portion that binds to an antigen and, optionally, a scaffold or framework portion that allows the antigen binding portion to adopt a conformation that promotes binding of the antigen binding protein to the antigen. Examples of antigen binding proteins include antibodies, antibody fragments (e.g., an antigen binding portion of an antibody), antibody derivatives, and antibody analogs. The antigen binding protein can comprise an alternative protein scaffold or artificial scaffold with grafted CDRs or CDR derivatives. Such scaffolds include, but are not limited to, antibody-derived scaffolds comprising mutations introduced to, for example, stabilize the three-dimensional structure of the antigen binding protein as well as wholly synthetic scaffolds comprising, for example, a biocompatible polymer. See, for example, Korndorfer et al., *Proteins: Structure, Function, and Bioinformatics*, (2003) Volume 53, Issue 1:121-129; Roque et al., *Biotechnol. Prog.*, 2004, 20:639-654. In addition, peptide antibody mimetics ("PAMs") can be used, as well as scaffolds based on antibody mimetics utilizing fibronectin components as a scaffold.

Certain antigen binding proteins described herein are antibodies or are derived from antibodies. Such antigen binding proteins include, but are not limited to, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies, antibody mimetics, chimeric antibodies, humanized antibodies, human antibodies, antibody fusions, antibody conjugates, single chain antibodies, and fragments thereof, respectively. In some instances, the antigen binding protein is an immunological fragment of an antibody (e.g., a Fab, a Fab', a F(ab')₂, or a scFv). The various structures are further described and defined herein.

Certain antigen binding proteins that are provided may comprise one or more CDRs as described herein (e.g., 1, 2,

3, 4, 5, 6 or more CDRs). In some instances, the antigen binding protein comprises (a) a polypeptide structure and (b) one or more CDRs that are inserted into and/or joined to the polypeptide structure. The polypeptide structure can take a variety of different forms. For example, it can be, or comprise, the framework of a naturally occurring antibody, or fragment or variant thereof, or may be completely synthetic in nature. Examples of various polypeptide structures are further described below.

An antigen binding protein of the invention is said to “specifically bind” its target antigen when the dissociation equilibrium constant (KD) is $\leq 10^{-8}$ M. The antigen binding protein specifically binds antigen with “high affinity” when the KD is $\leq 5 \times 10^{-9}$ M, and with “very high affinity” when the KD is $\leq 5 \times 10^{-10}$ M. In one embodiment the antigen binding protein will bind to human IL-23 with a KD of $\leq 5 \times 10^{-12}$ M, and in yet another embodiment it will bind with a KD $\leq 5 \times 10^{-13}$ M. In another embodiment of the invention, the antigen binding protein has a KD of $\leq 5 \times 10^{-12}$ M and an K_{off} of about $\leq 5 \times 10^{-6}$ 1/s. In another embodiment, the K_{off} is $\leq 5 \times 10^{-7}$ 1/s.

Another aspect provides an antigen binding protein having a half-life of at least one day in vitro or in vivo (e.g., when administered to a human subject). In one embodiment, the antigen binding protein has a half-life of at least three days. In another embodiment, the antibody or portion thereof has a half-life of four days or longer. In another embodiment, the antibody or portion thereof has a half-life of eight days or longer. In another embodiment, the antibody or antigen binding portion thereof is derivatized or modified such that it has a longer half-life as compared to the underivatized or unmodified antibody. In another embodiment, the antigen binding protein contains point mutations to increase serum half life, such as described in WIPO Publication No. WO 00/09560.

In embodiments where the antigen binding protein is used for therapeutic applications, an antigen binding protein can reduce, inhibit, interfere with or modulate one or more biological activities of IL-23, such inducing production of proinflammatory cytokines. IL-23 has many distinct biological effects, which can be measured in many different assays in different cell types; examples of such assays and known and are provided herein.

Some of the antigen binding proteins that are provided have the structure typically associated with naturally occurring antibodies. The structural units of these antibodies typically comprise one or more tetramers, each composed of two identical couplets of polypeptide chains, though some species of mammals also produce antibodies having only a single heavy chain. In a typical antibody, each pair or couplet includes one full-length “light” chain (in certain embodiments, about 25 kDa) and one full-length “heavy” chain (in certain embodiments, about 50-70 kDa). Each individual immunoglobulin chain is composed of several “immunoglobulin domains”, each consisting of roughly 90 to 110 amino acids and expressing a characteristic folding pattern. These domains are the basic units of which antibody polypeptides are composed. The amino-terminal portion of each chain typically includes a variable region that is responsible for antigen recognition. The carboxy-terminal portion is more conserved evolutionarily than the other end of the chain and is referred to as the “constant region” or “C region”. Human light chains generally are classified as kappa and lambda light chains, and each of these contains one variable region and one constant domain (CL1).z Heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon chains, and these define the antibody’s isotype as

IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subtypes, including, but not limited to, IgG1, IgG2, IgG3, and IgG4. IgM subtypes include IgM, and IgM2. IgA subtypes include IgA1 and IgA2. In humans, the IgA and IgD isotypes contain four heavy chains and four light chains; the IgG and IgE isotypes contain two heavy chains and two light chains; and the IgM isotype contains five heavy chains and five light chains. The heavy chain constant region (CH) typically comprises one or more domains that may be responsible for effector function. The number of heavy chain constant region domains will depend on the isotype. IgG heavy chains, for example, each contains three CH region domains known as CH1, CH2 and CH3. The antibodies that are provided can have any of these isotypes and subtypes, for example, the IL-23 antigen binding protein is of the IgG1, IgG2, or IgG4 subtype. If an IgG4 is desired, it may also be desired to introduce a point mutation (CPSCP→CP-PCP) in the hinge region as described in Bloom et al., 1997, Protein Science 6:407) to alleviate a tendency to form intra-H chain disulfide bonds that can lead to heterogeneity in the IgG4 antibodies. Antibodies provided herein that are of one type can be changed to a different type using subclass switching methods. See, e.g., Lantto et al., 2002, Methods Mol. Biol. 178:303-316.

In full-length light and heavy chains, the variable and constant regions are joined by a “J” region of about twelve or more amino acids, with the heavy chain also including a “D” region of about ten more amino acids. See, e.g., Fundamental Immunology, 2nd ed., Ch. 7 (Paul, W., ed.) 1989, New York: Raven Press. The variable regions of each light/heavy chain pair typically form the antigen binding site.

Variable Regions

Various heavy chain and light chain variable regions (or domains) provided herein are depicted in TABLES 1 and 2. Each of these variable regions may be attached, for example, to heavy and light chain constant regions described above. Further, each of the so generated heavy and light chain sequences may be combined to form a complete antigen binding protein structure.

Provided are antigen binding proteins that contain at least one heavy chain variable region (VH) selected from the group consisting of VH1, VH2, VH3, VH4, VH5, VH6, VH7, VH8, VH9, VH10, VH11, VH12, VH13, VH14, VH15 and VH16 and/or at least one light chain variable region (VL) selected from the group consisting of VL1, VL2, VL3, VL4, VL5, VL6, VL7, VL8, VL9, VL10, VL11, VL12, VL13, VL14, VL15, and VL16 as shown in TABLES 1 and 2 below.

Each of the heavy chain variable regions listed in TABLE 2 may be combined with any of the light chain variable regions shown in TABLE 1 to form an antigen binding protein. In some instances, the antigen binding protein includes at least one heavy chain variable region and/or one light chain variable region from those listed in TABLES 1 and 2. In some instances, the antigen binding protein includes at least two different heavy chain variable regions and/or light chain variable regions from those listed in TABLES 1 and 2. The various combinations of heavy chain variable regions may be combined with any of the various combinations of light chain variable regions.

In other instances, the antigen binding protein contains two identical light chain variable regions and/or two identical heavy chain variable regions. As an example, the antigen binding protein may be an antibody or immunologically functional fragment that comprises two light chain variable regions and two heavy chain variable regions in

17

combinations of pairs of light chain variable regions and pairs of heavy chain variable regions as listed in TABLES 1 and 2. Examples of such antigen binding proteins comprising two identical heavy chain and light chain variable regions include: Antibody A VH14/VL14; Antibody B VH9/VL9; Antibody C VH10/VL10; Antibody D VH15/VL15; Antibody E VH1/VL1, Antibody F VH11/VL11; Antibody G VH12/VL12; Antibody H VH13/VL13; Antibody I VH8/VL8; Antibody J VH3/VL3; Antibody K VH7/VL7; Antibody L VH4/VL4; Antibody M VH5/VL5 and Antibody N VH6/VL6.

Some antigen binding proteins that are provided comprise a heavy chain variable region and/or a light chain variable region comprising a sequence of amino acids that differs from the sequence of a heavy chain variable region and/or a light chain variable region selected from TABLES 1 and 2 at only 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino

18

acid residues, wherein each such sequence difference is independently either a deletion, insertion or substitution of one amino acid. The light and heavy chain variable regions, in some antigen binding proteins, comprise sequences of amino acids that have at least 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity to the amino acid sequences provided in TABLES 1 and 2. Still other antigen binding proteins, e.g., antibodies or immunologically functional fragments, also include variant heavy chain region forms and/or variant light chain region forms as described herein.

The term "identity" refers to a relationship between the sequences of two or more polypeptide molecules or two or more polynucleotides, as determined by aligning and comparing the sequences. "Percent identity" means the percent of identical residues between the amino acids or nucleotides in the compared molecules and is calculated based on the size of the smallest of the molecules being compared.

TABLE 1

Exemplary Variant Light Chain Region Sequences						
FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4
V _L 1	QSVLTQPPSVSGAPQQRVTISCTGSSSNIGAGYDVHMYQQVPGTAPKLLIYGSCNRP	SGVDPDRP	SGSKGTSASLAITGLQAEDEADYYCQSYDSSLG	SNVFGG	GTKLTVL	
SEQ ID NO: 1						
V _L 2	QSVLTQPPSVSGAPQQRVTISCTGSSSNIGAGYDVHMYQQVPGTAPKLLIYGSCNRP	SGVDPDRP	SGSKGTSASLAITGLQAEDEADYYCQSYDSSLG	SNVFGG	GTKLTVL	
SEQ ID NO: 3						
V _L 3	QAVLTQPPSSLGASPCASASLTCTLRSGINVGTYRIYNYQKPGSPQYLLRYKSDSDKQCGSGVP	SRFSGSKDASANAGILLISGLQSEDEADYYCMIWHSSASV	FGG	GTKLTVL		
SEQ ID NO: 4						
V _L 4	QAVLTQPPSSLGASPCASASLTCTLRSGINVGTYRIYNYQKPGSPQYLLRYKSDSDKQCGSGVP	SRFSGSKDASANAGILLISGLQSEDEADYYCMIWHSSASV	FGG	GTKLTVL		
SEQ ID NO: 4						
V _L 5	QPVLTQPPSASASLGASVTLTCTLNSGYSDYKVDWYQQRPGKGRPFVYMRVGTGGIVGSKDGI	PDRPSVLGSLNRYLTIKNIQEEDESDYHCCADHSGSGSNFVYV	FGTGT	KVTVL		
SEQ ID NO: 7						
V _L 6	QPVLTQPPSASASLGASVTLTCTLSSGYSDYKVDWYQQRPGKGRPFVYMRVGTGGIVGSKDGI	PDRPSVLGSLNRYLTIKNIQEEDESDYHCCADHSGSGNNFVYV	FGTGT	KVTVL		
SEQ ID NO: 9						
V _L 7	QPELTQPPSASASLGASVTLTCTLSSGYSDYKVDWYQLRPGKGRPFVYMRVGTGGIVGSKDGI	PDRPSVLGSLNRSLTIKNIQEEDESDYHCCADHSGSGNNFVYV	FGTGT	KVTVL		
SEQ ID NO: 11						
V _L 8	DIQLTPSPSSVSASVGDRTVITCRASQGIACWLAWYQQKPKGAPKLLIYAASSLQSGVP	SRFSGSGSGTDFTLTISSLQPEDFATYYCQQADSFP	FTFGG	GTKVEIK		
SEQ ID NO: 13						
V _L 9	DIQMTQSPSSVSASVGDRTVITCRASQVVISSWLAWYQQKPKGAPKLLIYAASSLQSGVP	SRFSGSVSGTDFTLTISSLQPEDFATYYCQQANSFP	FTFGG	TKVDIK		
SEQ ID NO: 15						
V _L 10	DIQMTQSPSSVSASVGDRTVITCRASQSGSSWFAWYQQKPKGAPKLLIYAASSLQSGVP	SRFSGSGSGTDFTLTISSLQPEDFATYYCQQANSFP	FTFGG	TKVDIK		
SEQ ID NO: 17						
V _L 11	DSQMTQSPSSVSASVGDRTVITCRASQGISSWFAWYQQKQAPNLLIYAASSLQSGVP	SRFSGSGSGTEFTLTISSLQPEDFATYYCQQANSFP	FTFGG	TKVDIK		
SEQ ID NO: 19						
V _L 12	DIQMTQSPSSVSASVGDRTVITCRAGQVISSWLAWYQQKPKGAPKLLIYAASSLQSGVP	SRFSGSGSGTDFTLTISSLQPDDEFATYYCQQATSF	PLTFGG	GTKVEIK		
SEQ ID NO: 21						
V _L 13	DIQMTQSPSSVSASVGDRTVITCRASQGFSGWLAWYQQKPKGAPKLLIYAASSLQSGVP	SRFSGSGSGTDFTLTISSLQPEDFATYYCQQANSFP	FTFGG	TKVDIK		
SEQ ID NO: 23						
V _L 14	DIQLTPSPSSVSASVGDRTVITCRASQVVISSWFAWYQQKQAPNLLIYAASSLQSGVP	SRFSGSGSGTDFTLTISSLQPADFATYFCQQANSFP	FTFGG	TKVDIK		
SEQ ID NO: 25						

TABLE 1-continued

Exemplary Variant Light Chain Region Sequences						
FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4
V _L 15	DIQMTQSPSSVSASVGDRTITTCRASQGSSEWFAWYQQKPGKAPKLLIYAASSLQSGVPSRFSGSGTDFTLTISLQPEDFATYYCOQANSFPFTFGPGTKVDIK					
SEQ ID NO: 27						
V _L 16	DIQMTQSPSSLSASVGDRTITTCRASQGIKNDLGWYQQKPGKAPKRLIYAASSLQSGVPSRFSGSGTEFTLTISLQPEDFATYYCLOHNSYPPTFGQGTKVEIE					
SEQ ID NO: 29						

TABLE 2

Exemplary Variant Heavy Chain Region Sequences							
	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4
V _H 1	QVQLVESGGGVVQPGRSURLSCAASGFTFSSYCYGMHWVRQAPGKGLEWAVAIWYDGSNEYADSVKGRFTI SRDNSKNTLYLQMNLSRAEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 31						
V _H 2	QVQLVESGGGVVQPGRSURLSCAASGFTFSSYCYGMHWVRQAPGKGLEWAVAIWYDGSNKYADSVKGRFTI SRDNSKNTLYLQMNLSRAEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 33						
V _H 3	QVQLVESGGGVVQPGRSURLSCAASGFTFSSYCYGMHWVRQAPGKGLEWAVAVISFDGSLKIYADSVKGRFTI SRDNSKNTLYLQMNLSRAEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 34						
V _H 4	QVQLVESGGGVVQPGRSURLSCAASGFTFSSYAMHWVRQAPGKGLEWLSVISHDGSIKIYADSVKGRFTI SRDNSKNTLYLQMNLSRAEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 36						
V _H 5	EVQLVESGGGLVQPGGSURLSCAASGFTFSSYSMMNWVRQAPGKGLEWVSYYISRSSTIYIADSVKGRFTI SRDNAKNSLYLQMNLSRDEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 38						
V _H 6	EVQLVESGGGLVQPGGSURLSCAASGFTFSTYSMMNWVRQAPGKGLEWVSYYISSSSSTRYHADSVKGRFTI SRDNAKNSLYLQMNLSRDEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 40						
V _H 7	EVQLVESGGGLVQPGGSURLSCVVS GFTFSSFSMMNWVRQAPGKGLEWVSYYISRSSTIYIADSVKGRFTI SRDNAKNSLYLQMNLSRDEDTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 42						
V _H 8	QVQLQESGPGLVKPSETLSLTCTVSGGSI STYYNSWNRQAPGKGLEWIGLIYTSGSTNYNP SLKSRVTISVDTSKNQPSLRLTSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 44						
V _H 9	QVQLQESGPGLVKPSETLSLTCTVSGGSI SSGGY YNSWNRQHPGKGLEWIGCHIHYSGNTYNP SLKSRVTISVDTSKNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 46						
V _H 10	QVQLQESGPGLVKPSETLSLTCTVSGGSI SSGGY YNSWNRQHPGKGLEWIGYIYYSGSSYNP SLKSRVTISVDTSQNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 48						
V _H 11	QVQLQESGPGLVKPSETLSLTCTVSGGSI SSGGY YNSWNRQHPGKGLEWIGYIYYSGSTYNP SLKSRVTISVDTSKNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 50						
V _H 12	QVQLQESGPRLVKPSETLSLTCTVSGDSI SSY FNSWNRQPPGKGLEWIGYIYYSGSTYNP SLKSRVTI SIDTSKNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 52						
V _H 13	QVQLQESGPGLVKPSETLSLTCTVSGGSI SSGGY YNSWNRQHPGKGLEWIGYIYYSGNTYNP SLKSRVTISVDTSKNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 54						
V _H 14	QVQLQESGPGLVKPSETLSLTCTVSGGSI SSGGY YNSWNRQHPGKGLEWIGYIYYSGSTYNP SLKSRVIMSDTSKNQPSLKLSSVTAADTAVYYC						ADFNWGQGTMTVTSS
	SEQ ID NO: 56						

TABLE 2-continued

Exemplary Variant Heavy Chain Region Sequences							
	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4
V _H 15	QVQLQESGPGLVKPSQTLTCTVSGGSINS GGYYW SWIRQHPGKGLEWIGI YYSGSSYYNP SLKSRVTISVDTSKNQFSLKLSVTAADTAVYYCAP DRGHYYGMDV WGQGTITVTVSS						
SEQ ID NO: 58							
V _H 16	QVQLVESGGGVVQPGRSLRLSCAASGFTFSS YGMH WVRQAPGKGLEWV ALLIWIYDGSNKYADSVKGR FTISRDN SKNTLYLQMNSLR AEDTAVYYC ARENTVTIIYYNYGMDV WGQGTITVTVSS						
SEQ ID NO: 60							

For these calculations, gaps in alignments (if any) must be addressed by a particular mathematical model or computer program (i.e., an "algorithm"). Methods that can be used to calculate the identity of the aligned nucleic acids or polypeptides include those described in *Computational Molecular Biology*, (Lesk, A. M., ed.), 1988, New York: Oxford University Press; *Biocomputing Informatics and Genome Projects*, (Smith, D. W., ed.), 1993, New York: Academic Press; *Computer Analysis of Sequence Data, Part I*, (Griffin, A. M., and Griffin, H. G., eds.), 1994, New Jersey: Humana Press; von Heinje, G., 1987, *Sequence Analysis in Molecular Biology*, New York: Academic Press; *Sequence Analysis Primer*, (Gribskov, M. and Devereux, J., eds.), 1991, New York: M. Stockton Press; and Carillo et al., 1988, *SIAM J. Applied Math.* 48:1073.

In calculating percent identity, the sequences being compared are aligned in a way that gives the largest match between the sequences. The computer program used to determine percent identity is the GCG program package, which includes GAP (Devereux et al., 1984, *Nucl. Acid Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, Wis.). The computer algorithm GAP is used to align the two polypeptides or polynucleotides for which the percent sequence identity is to be determined. The sequences are aligned for optimal matching of their respective amino acid or nucleotide (the "matched span", as determined by the algorithm). A gap opening penalty (which is calculated as 3× the average diagonal, wherein the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. In certain embodiments, a standard comparison matrix (see, Dayhoff et al., 1978, *Atlas of Protein Sequence and Structure* 5:345-352 for the PAM 250 comparison matrix; Henikoff et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89:10915-10919 for the BLOSUM 62 comparison matrix) is also used by the algorithm.

Recommended parameters for determining percent identity for polypeptides or nucleotide sequences using the GAP program are the following: Algorithm: Needleman et al., 1970, *J. Mol. Biol.* 48:443-453; Comparison matrix: BLOSUM 62 from Henikoff et al., 1992, supra; Gap Penalty: 12 (but with no penalty for end gaps), Gap Length Penalty: 4, Threshold of Similarity: 0. Certain alignment schemes for aligning two amino acid sequences may result in matching of only a short region of the two sequences and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, the selected alignment method (GAP program) can be adjusted if so desired to result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

The heavy and light chain variable regions disclosed herein include consensus sequences derived from groups of related antigen binding proteins. The amino acid sequences of the heavy and light chain variable regions were analyzed for similarities. Four groups emerged, one group having kappa light chain variable regions, (V_H9/V_L9, V_H10/V_L10, V_H11/V_L11, V_H13/V_L13, V_H14/V_L14 and V_H15/V_L15) and three groups having lambda light chain variable regions: lambda group 1 (V_H5/V_L5, V_H6/V_L6 and V_H7/V_L7), lambda group 2 (V_H3/V_L3 and V_H4/V_L4), and lambda group 3 (V_H1/V_L1 and V_H2/V_L2). Light chain germlines represented include VK1/A30 and VK1/L19. Light chain lambda germ-

lines represented include VL1/1e, VL3/3p, VL5/5c and VL9/9a. Heavy chain germlines represented include VH3/3-30, VH3/3-30.3, VH3/3-33, VH3/3-48, VH4/4-31 and VH4/4-59. As used herein, a "consensus sequence" refers to amino acid sequences having conserved amino acids common among a number of sequences and variable amino acids that vary within given amino acid sequences. Consensus sequences may be determined using standard phylogenetic analyses of the light and heavy chain variable regions corresponding to the IL-23 antigen binding proteins disclosed herein.

The light chain variable region consensus sequence for the kappa group is DX₁QX₂TQSPSSVSA SVGDRVTITCRASQGX₃X₄SX₅WX₆AWYQKPGX₇ 15 APX₈LLIYAASSLQSGVPSR FS GSX₉SGTX₁₀ FTLTISSLQPX₁₁DFATYX₁₂CQQANSFPFTFGPGTKV DX₁₃K (SEQ ID NO:30) where X₁ is selected from I or S; X₂ is selected from M or L; X₃ is selected from G or V and X₄ is selected from S, F or I; X₅ is selected from S or G; X₆ is selected from F or L; X₇ is selected from K or Q; X₈ is selected from K, N or S; X₉ is selected from G or V; X₁₀ is selected from D or E, X₁₁ is selected from E or A; X₁₂ is selected from Y or F; and X₁₃ is selected from I, V or F.

The light chain variable region consensus sequence for lambda group 1 is QPX₁LTQPPSASASLGA SVTLTCTLX₂SGYSDYKVDWYQX₃RPGKGPFRVMR VGTGGX₄VGSKGX₅GIPDRFSVLGSLNRX₆ 25 LTIKNIQEEDSDYHCGADHGSGX₇NFVYVFGTGTK VTVL (SEQ ID NO:61) where X₁ is selected from V or E; X₂ is selected from N or S; X₃ is selected from Q or L and X₄ is selected from I or T; X₅ is selected from D or E; X₆ is selected from Y or S; and X₇ is selected from S or N.

The light chain variable region consensus sequence for lambda group 3 is QSVLTQPPSVSGAPGQ RVTISCTGSSSNX₁GAGYDVHWYQQX₂PGTAPKLLIY GSX₃NRPSGVPDRFSGSKSGTSASLAITGLQAEDEADY YCQSYDSSLSGWVFGGGTX₄RLTVL (SEQ ID NO:139) 35 where X₁ is selected from T or I; X₂ is selected from V or L; X₃ is selected from G or N and X₄ is selected from R or K.

The heavy chain variable region consensus sequence for the kappa group is QVQLQESGPGLVKPSQ LSLT CTVSGGSIX₁SGGYWX₂WIRQHPGKGLEWIGX₃IX₄ YSGX₅X₆YYNP SLK SRX₇TX₈SVDT SX₉NQFSLX₁₀LSS 45 VTAADTAVYYCAX₁₁X₁₂RGX₁₃YYGMDVWGQGTTV TVSS (SEQ ID NO:140) where X₁ is selected from N or S; X₂ is selected from S or T; X₃ is selected from Y or H and X₄ is selected from Y or H; X₅ is selected from S or N; X₆ is selected from S or T; X₇ is selected from V or I; X₈ is selected from I or M; X₉ is selected from K or Q; X₁₀ is selected from K or S, X₁₁ is selected from R or K; X₁₂ is selected from D or N; and X₁₃ is selected from H, F or Y.

The heavy chain variable region consensus sequence for lambda group 1 is EVQLVESGGGLVQPGGSL RLSCX₁X₂SGFTFSX₃X₄SMNWVRQAPGKGLEWVS Y ISSX₅SSTX₆YX₇ADSVKGRFTISRDN AKN SLYLQMNS 55 LRDEDTAVYYCARRIAAGX₈X₉X₁₀YYYAX₁₁DVW GQGT TVTVSS (SEQ ID NO:141) where X₁ is selected from A or V; X₂ is selected from A or V; X₃ is selected from T or S and X₄ is selected from Y or F; X₅ is selected from S or R; X₆ is selected from R or I; X₇ is selected from H, Y or I; X₈ is selected from P or G; X₉ is selected from W or F; X₁₀ is selected from G or H and X₁₁ is selected from M or L.

The heavy chain variable region consensus sequence for lambda group 2 is QVQLVESGGGVVQPGRSRLRLSC AASGFTFSYX₁MHWVRQAPGKGLEWX₂X₃VISX₄

DGSX₃KYYAD SV KGRFTISRDN SKNTLYLQMNSL-
RAEDTAVYYCARERTITLSGSYFDYWGQGLTVTVSS
(SEQ ID NO:142) where X₁ is selected from G or A; X₂ is
selected from V or L; X₃ is selected from A or S and X₄ is
selected from F or H and X₅ is selected from L or I.

The heavy chain variable region consensus sequence for
lambda group 3 is QVQLVESGGG VVQPGRLRLS-
CAASGFTSSYGMHWVRQAPGKGLEWVAVIW
YDGSNX₁YYADSV KG RFTISRDN SKNTLYLQMNS
LRAEDTAVYYCARDRGYX₂SSWYPDAFDIWGQGT
MVTVSS (SEQ ID NO: 143) where X₁ is selected from E
or K and X₂ is selected from T or S.

Complementarity Determining Regions

Complementarity determining regions or "CDRs" are
embedded within a framework in the heavy and light chain
variable regions where they constitute the regions respon-
sible for antigen binding and recognition. Variable domains
of immunoglobulin chains of the same species, for example,
generally exhibit a similar overall structure; comprising
relatively conserved framework regions (FR) joined by
hypervariable CDR regions. An antigen binding protein can
have 1, 2, 3, 4, 5, 6 or more CDRs. The variable regions
discussed above, for example, typically comprise three
CDRs. The CDRs from heavy chain variable regions and
light chain variable regions are typically aligned by the
framework regions to form a structure that binds specifically
on a target antigen (e.g., IL-23). From N-terminal to C-ter-
minal, naturally-occurring light and heavy chain variable
regions both typically conform to the following order of
these elements: FR1, CDR1, FR2, CDR2, FR3, CDR3 and
FR4. The CDR and FR regions of exemplary light chain
variable domains and heavy chain variable domains are
highlighted in TABLES 1 and 2. It is recognized that the
boundaries of the CDR and FR regions can vary from those
highlighted. Numbering systems have been devised for
assigning numbers to amino acids that occupy positions in
each of these domains. Complementarity determining
regions and framework regions of a given antigen binding
protein may be identified using these systems. Numbering

systems are defined in Kabat et al., Sequences of Proteins of
Immunological Interest, 5th Ed., US Dept. of Health and
Human Services, PHS, NIH, NIH Publication No. 91-3242,
1991, or Chothia & Lesk, 1987, *J. Mol. Biol.* 196:901-917;
Chothia et al., 1989, *Nature* 342:878-883. Other numbering
systems for the amino acids in immunoglobulin chains
include IMGT® (the international ImMunoGeneTics infor-
mation system; Lefranc et al, *Dev. Comp. Immunol.* 2005,
29:185-203); and AHO (Honegger and Pluckthun, *J. Mol.*
Biol. 2001, 309(3):657-670). The CDRs provided herein
may not only be used to define the antigen binding domain
of a traditional antibody structure, but may be embedded in
a variety of other polypeptide structures, as described herein.

The antigen binding proteins disclosed herein are poly-
peptides into which one or more CDRs may be grafted,
inserted, embedded and/or joined. An antigen binding pro-
tein can have, for example, one heavy chain CDR1
("CDRH1"), and/or one heavy chain CDR2 ("CDRH2"),
and/or one heavy chain CDR3 ("CDRH3"), and/or one light
chain CDR1 ("CDRL1"), and/or one light chain CDR2
("CDRL2"), and/or one light chain CDR3 ("CDRL3").
Some antigen binding proteins include both a CDRH3 and
a CDRL3. Specific embodiments generally utilize combi-
nations of CDRs that are non-repetitive, e.g., antigen bind-
ing proteins are generally not made with two CDRH2
regions in one variable heavy chain region, etc. Antigen
binding proteins may comprise one or more amino acid
sequences that are identical to or that differ from the
amino acid sequences of one or more of the CDRs presented
in TABLE 3 at only 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,
14 or 15 amino acid residues, wherein each such sequence
difference is independently either a deletion, insertion or
substitution of one amino acid. The CDRs in some antigen
binding proteins comprise sequences of amino acids that
have at least 80%, 85%, 90%, 91%, 92, 93%, 94%, 95%,
96%, 97%, 98%, or 99% sequence identity to CDRs
sequence listed in TABLE 3. In some antigen binding
proteins, the CDRs are embedded into a "framework"
region, which orients the CDR(s) such that the proper
antigen binding properties of the CDR(s) is achieved.

TABLE 3

Exemplary CDRH and CDRL Sequences		
Exemplary CDRL Sequences		
CDRL1	CDRL2	CDRL3
TGSSSNTGAGYDVH SEQ ID NO: 62	GSGN RPS SEQ ID NO: 63	QSYDSSLGSGV SEQ ID NO: 64
TGSSNIGAGYDVH SEQ ID NO: 65	GSGNNRPS SEQ ID NO: 66	MIWHSSASV SEQ ID NO: 67
TLRSGINVGTYRIY SEQ ID NO: 68	YKSDSDKQGS SEQ ID NO: 69	GADHGSGSNFVYV SEQ ID NO: 70
TLNSGYSDYKV SEQ ID NO: 71	VGTGGIVGSKGD SEQ ID NO: 72	GADHGSGNNFVYV SEQ ID NO: 73
TLSSGYSDYKV SEQ ID NO: 74	VGTGGIVGSKGE SEQ ID NO: 75	QQANSFPPT SEQ ID NO: 76
RASQGFSGWLA SEQ ID NO: 77	VGTGGTVGSKGE SEQ ID NO: 78	QQATSFPLT SEQ ID NO: 79
RASQVISSWLA SEQ ID NO: 80	AASSLQS SEQ ID NO: 81	QQADSFPPPT SEQ ID NO: 82
RASQVISSWFA SEQ ID NO: 83		LQHNSYPPT SEQ ID NO: 84

TABLE 3-continued

Exemplary CDRH and CDRL Sequences		
RASQSSSWFA		
SEQ ID NO: 85		
RASQGISSWFA		
SEQ ID NO: 86		
RAGQVISSWLA		
SEQ ID NO: 87		
RASQGIAGWLA		
SEQ ID NO: 88		
RASQGIRNDLG		
SEQ ID NO: 89		
Exemplary CDRH Sequences		
CDRH1	CDRH2	CDRH3
SYGMH	VIWYDGSNEYADSVKG	DRGYTSSWYPDAFDI
SEQ ID NO: 91	SEQ ID NO: 92	SEQ ID NO: 93
SYAMH	VIWYDGSNKYYADSVKG	DRGYSSSWYPDAFDI
SEQ ID NO: 94	SEQ ID NO: 95	SEQ ID NO: 96
TYSMN	VISFDGSLKYYADSVKG	ERTTSLGSYFDY
SEQ ID NO: 97	SEQ ID NO: 98	SEQ ID NO: 99
SYSMN	VISHDGSIKYYADSVKG	RIAAAGGFHYYYALDV
SEQ ID NO: 100	SEQ ID NO: 101	SEQ ID NO: 102
SFSMN	YISSRSSTIYIADSVKG	RIAAAGPWGYYYAMDV
SEQ ID NO: 103	SEQ ID NO: 104	SEQ ID NO: 105
SGGYYWT	YISSSSSTRYHADSVKG	NRGYYYGMDV
SEQ ID NO: 106	SEQ ID NO: 107	SEQ ID NO: 108
SGGYYWS	YISSRSSTIYYADSVKG	NRGFYYGMDV
SEQ ID NO: 109	SEQ ID NO: 110	SEQ ID NO: 111
SYFWS	YIYYSGNTYYNPSLKS	DRGHYYGMDV
SEQ ID NO: 112	SEQ ID NO: 113	SEQ ID NO: 114
TTYWS	HIHYSGNTYYNPSLKS	DRGSYYGSDY
SEQ ID NO: 115	SEQ ID NO: 116	SEQ ID NO: 117
	YIYYSGSTYYNPSLKS	DRGYYYGVDV
	SEQ ID NO: 118	SEQ ID NO: 119
	YIYYSGSSYYNPSLKS	ENTVTIYYNYGMDV
	SEQ ID NO: 120	SEQ ID NO: 6
	YIYYSGSTNYNPSLKS	
	SEQ ID NO: 121	
	LIYTSGSTNYNPSLKS	
	SEQ ID NO: 122	
	LIWYDGSNKYYADSVKG	
	SEQ ID NO: 90	

Provided herein are CDR1 regions comprising amino acid residues 23-34 of SEQ ID NOs: 7 and 11; amino acid residues 24-34 of SEQ ID NOs: 9, 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 23-36 of SEQ ID NOs: 1, 3 and 4; amino acid residues 31-35 of SEQ ID NOs: 31, 33, 34, 38, 40, 44, 52 and 60 and amino acid residues 31-37 or SEQ ID NOs: 46, 48, 50, 54, 56 and 58.

CDR2 regions are provided comprising amino acid residues 50-56 of SEQ ID NOs: 9, 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 50-61 of SEQ ID NOs: 7 and 11; amino acid residues 52-62 of SEQ ID NO: 4; amino acid residues 50-65 of SEQ ID NOs: 31, 33, 44 and 52; amino acid residues 50-66 of SEQ ID NOs: 36, 38, 40, 42 and 60;

amino acid residues 52-58 of SEQ ID NOs: 1 and 3 and amino acid residues 52-67 of SEQ ID NOs: 46, 48, 50, 54, 56 and 58.

CDR3 regions comprising amino acid residues 89-97 of SEQ ID NOs: 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 91-101 of SEQ ID NOs: 1 and 3; amino acid residues 94-106 of SEQ ID NOs: 7, 9 and 11; amino acid residues 98-107 of SEQ ID NOs: 44 and 52; amino acid residues 97-105 of SEQ ID NO: 4; amino acid residues 99-110 of SEQ ID NOs: 34 and 36; amino acid residues 99-112 of SEQ ID NO: 112; amino acid residues 99-113 of SEQ ID NOs: 31 and 33; amino acid residues 99-114 of SEQ ID NOs: 38, 40 and 42; amino acid residues 100-109 of SEQ

ID NOs: 46, 48, 54, 56 and 58; and amino acid residues 101-019 of SEQ ID NO: 50; are also provided.

The CDRs disclosed herein include consensus sequences derived from groups of related sequences. As described previously, four groups of variable region sequences were identified, a kappa group and three lambda groups. The CDRL1 consensus sequence from the kappa group consists of RASQX₁X₂SX₃WX₄A (SEQ ID NO:123) where X₁ is selected from G or V; X₂ is selected from I, F or S; X₃ is selected from S or G and X₄ is selected from F or L. The CDRL1 consensus sequence from lambda group 1 consists of TLX₁SGYSDYKVD (SEQ ID NO:124) wherein X₁ is selected from N or S. The CDRL1 consensus sequences from lambda group 3 consists of TGSSSNX₁GAGYDVH (SEQ ID NO:125) wherein X₁ is selected from I or T.

The CDRL2 consensus sequence from lambda group 1 consists of VGTGGX₁VGSKGX₂ (SEQ ID NO: 126) wherein X₁ is selected from I or T and X₂ is selected from D or E. The CDRL2 consensus sequence from lambda group 3 consists of GSX₁NRPS (SEQ ID NO:127) wherein X₁ is selected from N or G.

The CDRL3 consensus sequences include GADHGSX₁NFVYV (SEQ ID NO:128) wherein X₁ is S or N.

The CDRH1 consensus sequence from the kappa group consists of SGGYYWX₁ (SEQ ID NO:129) wherein X₁ is selected from S or T. The CDRH1 consensus sequence from lambda group 1 consists of X₁X₂SMN (SEQ ID NO:131) wherein X₁ is selected from S or T and X₂ is selected from Y or F. The CDRH1 consensus sequence from lambda group 2 consists of SYX₁MH (SEQ ID NO:130), wherein X₁ is selected from G or A.

The CDRH2 consensus sequence from the kappa group consists of X₁IX₂YSGX₃X₄YYNPSLKS (SEQ ID NO:132) wherein X₁ is selected from Y or H; X₂ is selected from Y or H; X₃ is selected from S or N and X₄ is selected from T or S. The consensus sequence from lambda group 1 consists of YISSX₁SSTX₂YX₃ADSVKVG (SEQ ID NO:134) wherein X₁ is selected from R or S, X₂ is selected from I or R, X₃ is selected from I, H or Y. The consensus sequence from lambda group 2 consists of VISX₁DGSX₂KYYADSVKVG (SEQ ID NO:133) wherein X₁ is F or H and X₂ is L or T. The CDRH2 consensus sequence from lambda group 3 consists of VIWYDGSNX₁YYADSVKVG (SEQ ID NO:135) wherein X₁ is selected from K or E.

The CDRH3 consensus sequence from the kappa group consists of X₁RGX₂YYGMDV (SEQ ID NO:136) wherein X₁ is selected from N or D and X₂ is selected from H, Y or F. The CDRH3 consensus sequence from lambda group 1 consists of RIAAAGX₁X₂X₃YYYAX₄DV (SEQ ID NO:137) wherein X₁ is selected from G or P; X₂ is selected from F or W; X₃ is selected from H or G and X₄ is selected from L and M. The CDRH3 consensus sequence from lambda group 3 consists of DRGYX₁SSWYPDAFDI (SEQ ID NO:138) wherein X₁ is selected from S or T.

Monoclonal Antibodies

The antigen binding proteins that are provided include monoclonal antibodies that bind to IL-23. Monoclonal antibodies may be produced using any technique known in the art, e.g., by immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells can be immortalized using any technique known in the art, e.g., by fusing them with myeloma cells to produce hybridomas. Myeloma cells for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of

growing in certain selective media which support the growth of only the desired fused cells (hybridomas). Examples of suitable cell lines for use in mouse fusions include Sp-20, P3-X63/Ag8, P3-X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XXO Bul; examples of cell lines used in rat fusions include R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210. Other cell lines useful for cell fusions are U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6.

In some instances, a hybridoma cell line is produced by immunizing an animal (e.g., a transgenic animal having human immunoglobulin sequences) with an IL-23 immunogen; harvesting spleen cells from the immunized animal; fusing the harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; establishing hybridoma cell lines from the hybridoma cells, and identifying a hybridoma cell line that produces an antibody that binds an IL-23 polypeptide while sparing IL-12. Such hybridoma cell lines, and anti-IL-23 monoclonal antibodies produced by them, are aspects of the present application.

Monoclonal antibodies secreted by a hybridoma cell line can be purified using any technique known in the art. Hybridomas or mAbs may be further screened to identify mAbs with particular properties, such as the ability to inhibit IL-23-induced activity.

Chimeric and Humanized Antibodies

Chimeric and humanized antibodies based upon the foregoing sequences are also provided. Monoclonal antibodies for use as therapeutic agents may be modified in various ways prior to use. One example is a chimeric antibody, which is an antibody composed of protein segments from different antibodies that are covalently joined to produce functional immunoglobulin light or heavy chains or immunologically functional portions thereof. Generally, a portion of the heavy chain and/or light chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass. For methods relating to chimeric antibodies, see, for example, U.S. Pat. No. 4,816,567; and Morrison et al., 1985, *Proc. Natl. Acad. Sci. USA* 81:6851-6855. CDR grafting is described, for example, in U.S. Pat. Nos. 6,180,370, 5,693,762, 5,693,761, 5,585,089, and 5,530,101.

One useful type of chimeric antibody is a "humanized" antibody. Generally, a humanized antibody is produced from a monoclonal antibody raised initially in a non-human animal. Certain amino acid residues in this monoclonal antibody, typically from non-antigen recognizing portions of the antibody, are modified to be homologous to corresponding residues in a human antibody of corresponding isotype. Humanization can be performed, for example, using various methods by substituting at least a portion of a rodent variable region for the corresponding regions of a human antibody (see, e.g., U.S. Pat. No. 5,585,089, and No. 5,693,762; Jones et al., 1986, *Nature* 321:522-525; Riechmann et al., 1988, *Nature* 332:323-27; Verhoeyen et al., 1988, *Science* 239: 1534-1536).

In certain embodiments, constant regions from species other than human can be used along with the human variable region(s) to produce hybrid antibodies.

Fully Human Antibodies

Fully human antibodies are also provided. Methods are available for making fully human antibodies specific for a given antigen without exposing human beings to the antigen

("fully human antibodies"). One specific means provided for implementing the production of fully human antibodies is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated is one means of producing fully human monoclonal antibodies (mAbs) in mouse, an animal that can be immunized with any desirable antigen. Using fully human antibodies can minimize the immunogenic and allergic responses that can sometimes be caused by administering mouse or mouse-derivatized mAbs to humans as therapeutic agents.

Fully human antibodies can be produced by immunizing transgenic animals (usually mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. Antigens for this purpose typically have six or more contiguous amino acids, and optionally are conjugated to a carrier, such as a hapten. See, e.g., Jakobovits et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:2551-2555; Jakobovits et al., 1993, *Nature* 362:255-258; and Bruggermann et al., 1993, *Year in Immunol.* 7:33. In one example of such a method, transgenic animals are produced by inactivating the endogenous mouse immunoglobulin loci encoding the mouse heavy and light immunoglobulin chains therein, and inserting into the mouse genome large fragments of human genome DNA containing loci that encode human heavy and light chain proteins. Partially modified animals, which have less than the full complement of human immunoglobulin loci, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies that are immunospecific for the immunogen but have human rather than murine amino acid sequences, including the variable regions. For further details of such methods, see, for example, WIPO patent publications WO96/33735 and WO94/02602. Additional methods relating to transgenic mice for making human antibodies are described in U.S. Pat. Nos. 5,545,807; 6,713,610; 6,673,986; 6,162,963; 5,545,807; 6,300,129; 6,255,458; 5,877,397; 5,874,299 and 5,545,806; in WIPO patent publications WO91/10741, WO90/04036, and in EP 546073B1 and EP 546073A1.

The transgenic mice described above contain a human immunoglobulin gene minilocus that encodes unrearranged human heavy ([mu] and [gamma]) and [kappa] light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous [mu] and [kappa] chain loci (Lonberg et al., 1994, *Nature* 368:856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or [kappa] and in response to immunization, and the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgG [kappa] monoclonal antibodies (Lonberg et al., supra.; Lonberg and Huszar, 1995, *Intern. Rev. Immunol.* 13: 65-93; Harding and Lonberg, 1995, *Ann. N.Y. Acad. Sci.* 764:536-546). The preparation of such mice is described in detail in Taylor et al., 1992, *Nucleic Acids Research* 20:6287-6295; Chen et al., 1993, *International Immunology* 5:647-656; Tuaillon et al., 1994, *J. Immunol.* 152:2912-2920; Lonberg et al., 1994, *Nature* 368:856-859; Lonberg, 1994, *Handbook of Exp. Pharmacology* 113:49-101; Taylor et al., 1994, *International Immunology* 6:579-591; Lonberg and Huszar, 1995, *Intern. Rev. Immunol.* 13:65-93; Harding and Lonberg, 1995, *Ann. N.Y. Acad. Sci.* 764:536-546; Fishwild et al., 1996, *Nature Biotechnology* 14:845-85. See, further U.S. Pat. No. 5,545,806; No. 5,569,825; No. 5,625,126; No. 5,633,425; No. 5,789,650; No. 5,877,397; No. 5,661,016; No. 5,814,318; No. 5,874,299; and No. 5,770,

429; as well as U.S. Pat. No. 5,545,807; WIPO Publication Nos. WO 93/1227; WO 92/22646; and WO 92/03918. Technologies utilized for producing human antibodies in these transgenic mice are disclosed also in WIPO Publication No. WO 98/24893, and Mendez et al., 1997, *Nature Genetics* 15:146-156. For example, the HCo7 and HCo12 transgenic mice strains can be used to generate anti-IL-23 antibodies.

Using hybridoma technology, antigen-specific human mAbs with the desired specificity can be produced and selected from the transgenic mice such as those described above. Such antibodies may be cloned and expressed using a suitable vector and host cell, or the antibodies can be harvested from cultured hybridoma cells.

Fully human antibodies can also be derived from phage-display libraries (such as disclosed in Hoogenboom et al., 1991, *J. Mol. Biol.* 227:381; Marks et al., 1991, *J. Mol. Biol.* 222:581; WIPO Publication No. WO 99/10494). Phage display techniques mimic immune selection through the display of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice.

Bispecific or Bifunctional Antigen Binding Proteins

A "bispecific," "dual-specific" or "bifunctional" antigen binding protein or antibody is a hybrid antigen binding protein or antibody, respectively, having two different antigen binding sites, such as one or more CDRs or one or more variable regions as described above. In some instances they are an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Multispecific antigen binding protein or "multispecific antibody" is one that targets more than one antigen or epitope. Bispecific antigen binding proteins and antibodies are a species of multispecific antigen binding protein antibody and may be produced by a variety of methods including, but not limited to, fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai and Lachmann, 1990, *Clin. Exp. Immunol.* 79:315-321; Kostelny et al., 1992, *J. Immunol.* 148:1547-1553.

Immunological Fragments

Antigen binding proteins also include immunological fragments of an antibody (e.g., a Fab, a Fab', a F(ab')₂, or a scFv). A "Fab fragment" is comprised one light chain (the light chain variable region (V_L) and its corresponding constant domain (C_L)) and one heavy chain (the heavy chain variable region (V_H) and first constant domain (C_{H1})). The heavy chain of a Fab molecule cannot form a disulfide bond with another heavy chain molecule. A "Fab' fragment" contains one light chain and a portion of one heavy chain that also contains the region between the C_{H1} and C_{H2} domains, such that an interchain disulfide bond can be formed between the two heavy chains of two Fab' fragments to form an F(ab')₂ molecule. A "F(ab')₂ fragment" thus is composed of two Fab' fragments that are held together by a disulfide bond between the two heavy chains. A "Fv fragment" consists of the variable light chain region and variable heavy chain region of a single arm of an antibody. Single-chain antibodies "scFv" are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain, which forms an antigen binding region. Single chain antibodies are discussed in detail in WIPO Publication No. WO 88/01649, U.S. Pat. No. 4,946,778 and No. 5,260,203; Bird, 1988, *Science* 242:423; Huston et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879; Ward et al., 1989, *Nature* 334:544; de Graaf et al., 2002, *Methods Mol Biol.* 178:379-387; Kortt et al., 1997, *Prot. Eng.* 10:423; Kortt et al., 2001, *Biomol. Eng.*

18:95-108 and Kriangkum et al., 2001, *Biomol. Eng.* 18:31-40. A "Fc" region contains two heavy chain fragments comprising the C_H1 and C_H2 domains of an antibody. The two heavy chain fragments are held together by two or more disulfide bonds and by hydrophobic interactions of the C_H3 domains.

Also included are domain antibodies, immunologically functional immunoglobulin fragments containing only the variable region of a heavy chain or the variable region of a light chain. In some instances, two or more V_H regions are covalently joined with a peptide linker to create a bivalent domain antibody. The two V_H regions of a bivalent domain antibody may target the same or different antigens. Diabodies are bivalent antibodies comprising two polypeptide chains, wherein each polypeptide chain comprises V_H and V_L domains joined by a linker that is too short to allow for pairing between two domains on the same chain, thus allowing each domain to pair with a complementary domain on another polypeptide chain (see, e.g., Holliger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-48, 1993 and Poljak et al., *Structure* 2:1121-23, 1994). Similarly, tribodies and tetrabodies are antibodies comprising three and four polypeptide chains, respectively, and forming three and four antigen binding sites, respectively, which can be the same or different. Maxibodies comprise bivalent scFvs covalently attached to the Fc region of IgG₁, (see, e.g., Fredericks et al., 2004, *Protein Engineering, Design & Selection*, 17:95-106; Powers et al., 2001, *Journal of Immunological Methods*, 251:123-135; Shu et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:7995-7999; Hayden et al., 1994, *Therapeutic Immunology* 1:3-15).

Various Other Forms

Also provided are variant forms of the antigen binding proteins disclosed above, some of the antigen binding proteins having, for example, one or more conservative amino acid substitutions in one or more of the heavy or light chains, variable regions or CDRs listed in TABLES 1 and 2.

Naturally-occurring amino acids may be divided into classes based on common side chain properties: hydrophobic (norleucine, Met, Ala, Val, Leu, Ile); neutral hydrophilic (Cys, Ser, Thr, Asn, Gln); acidic (Asp, Glu); basic (His, Lys, Arg); residues that influence chain orientation (Gly, Pro); and aromatic (Trp, Tyr, Phe).

Conservative amino acid substitutions may involve exchange of a member of one of these classes with another member of the same class. Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and other reversed or inverted forms of amino acid moieties. Such substantial modifications in the functional and/or biochemical characteristics of the antigen binding proteins described herein may be achieved by creating substitutions in the amino acid sequence of the heavy and light chains that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulkiness of the side chain.

Non-conservative substitutions may involve the exchange of a member of one of the above classes for a member from another class. Such substituted residues may be introduced into regions of the antibody that are homologous with human antibodies, or into the non-homologous regions of the molecule.

In making such changes, according to certain embodiments, the hydropathic index of amino acids may be considered. The hydropathic profile of a protein is calculated by assigning each amino acid a numerical value ("hydropathy index") and then repetitively averaging these values along the peptide chain. Each amino acid has been assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics. They are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic profile in conferring interactive biological function on a protein is understood in the art (see, e.g., Kyte et al., 1982, *J. Mol. Biol.* 157:105-131). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, in certain embodiments, the substitution of amino acids whose hydropathic indices are within ± 2 is included. In some aspects, those which are within ± 1 are included, and in other aspects, those within ± 0.5 are included.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity, particularly where the biologically functional protein or peptide thereby created is intended for use in immunological embodiments, as in the present case. In certain embodiments, the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigen binding or immunogenicity, that is, with a biological property of the protein.

The following hydrophilicity values have been assigned to these amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 $\pm$ 1); glutamate (+3.0 $\pm$ 1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5 $\pm$ 1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5) and tryptophan (-3.4). In making changes based upon similar hydrophilicity values, in certain embodiments, the substitution of amino acids whose hydrophilicity values are within ± 2 is included, in other embodiments, those which are within ± 1 are included, and in still other embodiments, those within ± 0.5 are included. In some instances, one may also identify epitopes from primary amino acid sequences on the basis of hydrophilicity. These regions are also referred to as "epitopic core regions."

Exemplary conservative amino acid substitutions are set forth in TABLE 4.

TABLE 4

Conservative Amino Acid Substitutions							
Residue	Sub	Residue	Sub	Residue	Sub	Residue	Sub
Ala	Ser	Gln	Asn	Leu	Ile, Val	Thr	Ser
Arg	Lys	Glu	Asp	Lys	Arg, Gln, Glu	Trp	Tyr
Asn	Gln, His	Gly	Pro	Met	Leu, Ile	Tyr	Trp, Phe

TABLE 4-continued

Conservative Amino Acid Substitutions							
Residue	Sub	Residue	Sub	Residue	Sub	Residue	Sub
Asp	Glu	His	Asn, Gln	Phe	Met, Leu, Tyr	Val	Ile, Leu
Cys	Ser	Ile	Leu, Val	Ser	Thr	Thr	Ser

Residue = Original Residue
Sub = Exemplary Substitution

A skilled artisan will be able to determine suitable variants of polypeptides as set forth herein using well-known techniques. One skilled in the art may identify suitable areas of the molecule that may be changed without destroying activity by targeting regions not believed to be important for activity. The skilled artisan also will be able to identify residues and portions of the molecules that are conserved among similar polypeptides. In further embodiments, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in a protein that correspond to amino acid residues important for activity or structure in similar proteins. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues.

One skilled in the art can also analyze the 3-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of an antibody with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each desired amino acid residue. These variants can then be screened using assays for IL-23 activity, (see examples below) thus yielding information regarding which amino acids can be changed and which must not be changed. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acid positions where further substitutions should be avoided either alone or in combination with other mutations.

A number of scientific publications have been devoted to the prediction of secondary structure. See, Moulton, 1996, *Curr. Op. in Biotech.* 7:422-427; Chou et al., 1974, *Biochem.* 13:222-245; Chou et al., 1974, *Biochemistry* 113:211-222; Chou et al., 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47:45-148; Chou et al., 1979, *Ann. Rev. Biochem.* 47:251-276; and Chou et al., 1979, *Biophys. J.* 26:367-384. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins that have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural database (PDB) has

provided enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See, Holm et al., 1999, *Nucl. Acid. Res.* 27:244-247. It has been suggested (Brenner et al., 1997, *Curr. Op. Struct. Biol.* 7:369-376) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate.

Additional methods of predicting secondary structure include "threading" (Jones, 1997, *Curr. Opin. Struct. Biol.* 7:377-387; Sippl et al., 1996, *Structure* 4:15-19), "profile analysis" (Bowie et al., 1991, *Science* 253:164-170; Gribskov et al., 1990, *Meth. Enzym.* 183:146-159; Gribskov et al., 1987, *Proc. Nat. Acad. Sci.* 84:4355-4358), and "evolutionary linkage" (See, Holm, 1999, supra; and Brenner, 1997, supra).

In some embodiments, amino acid substitutions are made that: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter ligand or antigen binding affinities, and/or (4) confer or modify other physicochemical or functional properties on such polypeptides, such as maintaining the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation; maintaining or altering the charge or hydrophobicity of the molecule at the target site, or maintaining or altering the bulkiness of a side chain.

For example, single or multiple amino acid substitutions (in certain embodiments, conservative amino acid substitutions) may be made in the naturally-occurring sequence. Substitutions can be made in that portion of the antibody that lies outside the domain(s) forming intermolecular contacts). In such embodiments, conservative amino acid substitutions can be used that do not substantially change the structural characteristics of the parent sequence (e.g., one or more replacement amino acids that do not disrupt the secondary structure that characterizes the parent or native antigen binding protein). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed.), 1984, W. H. New York: Freeman and Company; *Introduction to Protein Structure* (Branden and Tooze, eds.), 1991, New York: Garland Publishing; and Thornton et al., 1991, *Nature* 354:105.

Additional variants include cysteine variants wherein one or more cysteine residues in the parent or native amino acid sequence are deleted from or substituted with another amino acid (e.g., serine). Cysteine variants are useful, inter alia when antibodies (for example) must be refolded into a biologically active conformation. Cysteine variants may have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

The heavy and light chain variable region and CDRs that are disclosed can be used to prepare antigen binding proteins that contain an antigen binding region that can specifically bind to an IL-23 polypeptide. "Antigen binding region" means a protein, or a portion of a protein, that specifically binds a specified antigen, such as the region that contains the amino acid residues that interact with an antigen and confer on the antigen binding protein its specificity and affinity for the target antigen. An antigen binding region may include one or more CDRs and certain antigen binding regions also include one or more "framework" regions. For example, one or more of the CDRs listed in TABLE 3 can be incorporated into a molecule (e.g., a polypeptide) covalently or nonco-

valently to make an immunoadhesion. An immunoadhesion may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDR(s) enable the immunoadhesion to bind specifically to a particular antigen of interest (e.g., an IL-23 polypeptide).

Other antigen binding proteins include mimetics (e.g., "peptide mimetics" or "peptidomimetics") based upon the variable regions and CDRs that are described herein. These analogs can be peptides, non-peptides or combinations of peptide and non-peptide regions. Fauchere, 1986, *Adv. Drug Res.* 15:29; Veber and Freidinger, 1985, *TINS* p. 392; and Evans et al., 1987, *J. Med. Chem.* 30:1229. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce a similar therapeutic or prophylactic effect. Such compounds are often developed with the aid of computerized molecular modeling. Generally, peptidomimetics are proteins that are structurally similar to an antigen binding protein displaying a desired biological activity, such as the ability to bind IL-23, but peptidomimetics have one or more peptide linkages optionally replaced by a linkage selected from, for example: $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{S}-$, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}-\text{CH}-$ (cis and trans), $-\text{COCH}_2-$, $-\text{CH}(\text{OH})\text{CH}_2-$, and $-\text{CH}_2\text{SO}-$, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used in certain embodiments to generate more stable proteins. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch, 1992, *Ann. Rev. Biochem.* 61:387), for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

Derivatives of the antigen binding proteins that are described herein are also provided. The derivatized antigen binding proteins can comprise any molecule or substance that imparts a desired property to the antigen binding protein or fragment, such as increased half-life in a particular use. The derivatized antigen binding protein can comprise, for example, a detectable (or labeling) moiety (e.g., a radioactive, colorimetric, antigenic or enzymatic molecule, a detectable bead (such as a magnetic or electrodense (e.g., gold) bead), or a molecule that binds to another molecule (e.g., biotin or Streptavidin)), a therapeutic or diagnostic moiety (e.g., a radioactive, cytotoxic, or pharmaceutically active moiety), or a molecule that increases the suitability of the antigen binding protein for a particular use (e.g., administration to a subject, such as a human subject, or other in vivo or in vitro uses). Examples of molecules that can be used to derivatize an antigen binding protein include albumin (e.g., human serum albumin) and polyethylene glycol (PEG). Albumin-linked and PEGylated derivatives of antigen binding proteins can be prepared using techniques well known in the art. In one embodiment, the antigen binding protein is conjugated or otherwise linked to transthyretin (TTR) or a TTR variant. The TTR or TTR variant can be chemically modified with, for example, a chemical selected from the group consisting of dextran, poly(n-vinyl pyrrolidone), polyethylene glycols, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols and polyvinyl alcohols.

Other derivatives include covalent or aggregative conjugates of IL-23 antigen binding proteins with other proteins or polypeptides, such as by expression of recombinant

fusion proteins comprising heterologous polypeptides fused to the N-terminus or C-terminus of an IL-23 antigen binding protein. For example, the conjugated peptide may be a heterologous signal (or leader) polypeptide, e.g., the yeast alpha-factor leader, or a peptide such as an epitope tag. IL-23 antigen binding protein-containing fusion proteins can comprise peptides added to facilitate purification or identification of the IL-23 antigen binding protein (e.g., poly-His). An IL-23 antigen binding protein also can be linked to the FLAG peptide as described in Hopp et al., 1988, *Bio/Technology* 6:1204; and U.S. Pat. No. 5,011,912. The FLAG peptide is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody (mAb), enabling rapid assay and facile purification of expressed recombinant protein. Reagents useful for preparing fusion proteins in which the FLAG peptide is fused to a given polypeptide are commercially available (Sigma, St. Louis, Mo.).

Oligomers that contain one or more IL-23 antigen binding proteins may be employed as IL-23 antagonists. Oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers. Oligomers comprising two or more IL-23 antigen binding proteins are contemplated for use, with one example being a homodimer. Other oligomers include heterodimers, homotrimers, heterotrimers, homotetramers, heterotetramers, etc. Oligomers comprising multiple IL-23-binding proteins joined via covalent or non-covalent interactions between peptide moieties fused to the IL-23 antigen binding proteins, are also included. Such peptides may be peptide linkers (spacers), or peptides that have the property of promoting oligomerization. Among the suitable peptide linkers are those described in U.S. Pat. Nos. 4,751,180 and 4,935,233. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of IL-23 antigen binding proteins attached thereto. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in WIPO Publication No. WO 94/10308; Hoppe et al., 1994, *FEBS Letters* 344:191; and Fanslow et al., 1994, *Semin. Immunol.* 6:267-278. In one approach, recombinant fusion proteins comprising an IL-23 antigen binding protein fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric IL-23 antigen binding protein fragments or derivatives that form are recovered from the culture supernatant.

Such oligomers may comprise from two to four IL-23 antigen binding proteins. The IL-23 antigen binding protein moieties of the oligomer may be in any of the forms described above, e.g., variants or fragments. Preferably, the oligomers comprise IL-23 antigen binding proteins that have IL-23 binding activity. Oligomers may be prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:10535; Bym et al., 1990, *Nature* 344:677; and Hollenbaugh et al., 1992 "Construction of Immunoglobulin Fusion Proteins", in *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1-10.19.11.

Also included are dimers comprising two fusion proteins created by fusing an IL-23 antigen binding protein to the Fc region of an antibody. The dimer can be made by, for example, inserting a gene fusion encoding the fusion protein into an appropriate expression vector, expressing the gene fusion in host cells transformed with the recombinant

expression vector, and allowing the expressed fusion protein to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield the dimer. Such Fc polypeptides include native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization also are included. Fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns. One suitable Fc polypeptide, described in WIPO Publication No. WO 93/10151 and U.S. Pat. Nos. 5,426,048 and 5,262,522, is a single chain polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Pat. No. 5,457,035, and in Baum et al., 1994, *EMBO J.* 13:3992-4001. The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WIPO Publication No. WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

Glycosylation

The antigen binding protein may have a glycosylation pattern that is different or altered from that found in the native species. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antigen binding protein is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antigen binding protein amino acid sequence may be altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the antigen binding protein is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free

sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in PCT Publication No. WO 87/05330, and in Aplin and Wriston, 1981, *CRC Crit. Rev. Biochem.*, pp. 259-306.

Removal of carbohydrate moieties present on the starting antigen binding protein may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge et al., 1981, *Anal. Biochem.* 118:131. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

Hence, aspects include glycosylation variants of the antigen binding proteins wherein the number and/or type of glycosylation site(s) has been altered compared to the amino acid sequences of the parent polypeptide. In certain embodiments, antigen binding protein variants comprise a greater or a lesser number of N-linked glycosylation sites than the parent polypeptide. Substitutions that eliminate or alter this sequence will prevent addition of an N-linked carbohydrate chain present in the parent polypeptide. For example, the glycosylation can be reduced by the deletion of an Asn or by substituting the Asn with a different amino acid. Antibodies typically have a N-linked glycosylation site in the Fc region.

Labels And Effector Groups

Antigen binding proteins may comprise one or more labels. The term "label" or "labeling group" refers to any detectable label. In general, labels fall into a variety of classes, depending on the assay in which they are to be detected: a) isotopic labels, which may be radioactive or heavy isotopes; b) magnetic labels (e.g., magnetic particles); c) redox active moieties; d) optical dyes; enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase); e) biotinylated groups; and f) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.). In some embodiments, the labeling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labeling proteins are known in the art. Examples of suitable labeling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the labeling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance.

Various methods for labeling proteins are known in the art and may be used as is seen fit.

The term "effector group" means any group coupled to an antigen binding protein that acts as a cytotoxic agent. Examples for suitable effector groups are radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I). Other suitable groups include toxins, therapeutic groups, or chemotherapeutic groups. Examples of suitable groups include calicheamicin, auristatins, geldanamycin and maytansine. In some embodiments, the effector group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance.

Polynucleotides Encoding IL-23 Antigen Binding Proteins

Polynucleotides that encode the antigen binding proteins described herein, or portions thereof, are also provided, including polynucleotides encoding one or both chains of an antibody, or a fragment, derivative, mutein, or variant thereof, polynucleotides encoding heavy chain variable regions or only CDRs, polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide, anti-sense nucleic acids for inhibiting expression of a polynucleotide, and complementary sequences of the foregoing. The polynucleotides can be any length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 85, 95, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleic acids in length, including all values in between, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger polynucleotide, for example, a vector. The polynucleotides can be single-stranded or double-stranded and can comprise RNA and/or DNA nucleic acids and artificial variants thereof (e.g., peptide nucleic acids).

Polynucleotides encoding certain antigen binding proteins, or portions thereof (e.g., full length antibody, heavy or light chain, variable domain, or a CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3) may be isolated from B-cells of mice that have been immunized with IL-23 or an immunogenic fragment thereof. The polynucleotide may be isolated by conventional procedures such as polymerase chain reaction (PCR). Phage display is another example of a known technique whereby derivatives of antibodies and other antigen binding proteins may be prepared. In one approach, polypeptides that are components of an antigen binding protein of interest are expressed in any suitable recombinant expression system, and the expressed polypeptides are allowed to assemble to form antigen binding protein molecules. Phage display is also used to derive antigen binding proteins having different properties (i.e., varying affinities for the antigen to which they bind) via chain shuffling, see Marks et al., 1992, *BioTechnology* 10:779.

Due to the degeneracy of the genetic code, each of the polypeptide sequences depicted herein are also encoded by a large number of other polynucleotide sequences besides those provided. For example, heavy chain variable domains provided herein in may be encoded by polynucleotide sequences SEQ ID NOs: 32, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, or 59. Light chain variable domains may be encoded by polynucleotide sequences SEQ ID NOs: 2, 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28. One of ordinary skill in the art will appreciate that the present application thus provides adequate written description and enablement for each degenerate nucleotide sequence encoding each antigen binding protein.

An aspect further provides polynucleotides that hybridize to other polynucleotide molecules under particular hybridization conditions. Methods for hybridizing nucleic acids, basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are well-known in the art. See, e.g., Sambrook, Fritsch, and Maniatis (2001, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. and *Current Protocols in Molecular Biology*, 1995, Ausubel et al., eds., John Wiley & Sons, Inc. As defined herein, a moderately stringent hybridization condition uses a prewashing solution containing 5× sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6×SSC, and a hybridization temperature of 55° C. (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42° C.), and washing conditions of 60° C., in 0.5×SSC, 0.1% SDS. A stringent hybridization condition hybridizes in 6×SSC at 45° C., followed by one or more washes in 0.1×SSC, 0.2% SDS at 68° C. Furthermore, one of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that polynucleotides comprising nucleic acid sequences that are at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to each other, including all values in between, typically remain hybridized to each other.

Changes can be introduced by mutation into a polynucleotide, thereby leading to changes in the amino acid sequence of a polypeptide (e.g., an antigen binding protein or antigen binding protein derivative) that it encodes. Mutations can be introduced using any technique known in the art, such as site-directed mutagenesis and random mutagenesis. Mutant polypeptides can be expressed and selected for a desired property. Mutations can be introduced into a polynucleotide without significantly altering the biological activity of a polypeptide that it encodes. For example, substitutions at non-essential amino acid residues. Alternatively, one or more mutations can be introduced into a polynucleotide that selectively change the biological activity of a polypeptide that it encodes. For example, the mutation can quantitatively or qualitatively change the biological activity, such as increasing, reducing or eliminating the activity and changing the antigen specificity of an antigen binding protein.

Another aspect provides polynucleotides that are suitable for use as primers or hybridization probes for the detection of nucleic acid sequences. A polynucleotide can comprise only a portion of a nucleic acid sequence encoding a full-length polypeptide, for example, a fragment that can be used as a probe or primer or a fragment encoding an active portion (e.g., an IL-23 binding portion) of a polypeptide. Probes based on the sequence of a nucleic acid can be used to detect the nucleic acid or similar nucleic acids, for example, transcripts encoding a polypeptide. The probe can comprise a label group, e.g., a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used to identify a cell that expresses the polypeptide.

Methods of Expressing Antigen Binding Proteins

The antigen binding proteins provided herein may be prepared by any of a number of conventional techniques. For example, IL-23 antigen binding proteins may be produced by recombinant expression systems, using any technique known in the art. See, e.g., *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet et al. (eds.) Plenum Press, New York (1980); and

Antibodies: A Laboratory Manual, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1988).

Expression systems and constructs in the form of plasmids, expression vectors, transcription or expression cassettes that comprise at least one polynucleotide as described above are also provided herein, as well host cells comprising such expression systems or constructs. As used herein, "vector" means any molecule or entity (e.g., nucleic acid, plasmid, bacteriophage or virus) suitable for use to transfer protein coding information into a host cell. Examples of vectors include, but are not limited to, plasmids, viral vectors, non-episomal mammalian vectors and expression vectors, for example, recombinant expression vectors. Expression vectors, such as recombinant expression vectors, are useful for transformation of a host cell and contain nucleic acid sequences that direct and/or control (in conjunction with the host cell) expression of one or more heterologous coding regions operatively linked thereto. An expression construct may include, but is not limited to, sequences that affect or control transcription, translation, and, if introns are present, affect RNA splicing of a coding region operably linked thereto. "Operably linked" means that the components to which the term is applied are in a relationship that allows them to carry out their inherent functions. For example, a control sequence, e.g., a promoter, in a vector that is "operably linked" to a protein coding sequence are arranged such that normal activity of the control sequence leads to transcription of the protein coding sequence resulting in recombinant expression of the encoded protein.

Another aspect provides host cells into which an expression vector, such as a recombinant expression vector, has been introduced. A host cell can be any prokaryotic cell (for example, *E. coli*) or eukaryotic cell (for example, yeast, insect, or mammalian cells (e.g., CHO cells)). Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably transfected with the introduced polynucleotide can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die), among other methods.

Antigen binding proteins can be expressed in hybridoma cell lines (e.g., in particular antibodies may be expressed in hybridomas) or in cell lines other than hybridomas. Expression constructs encoding the antigen binding proteins can be used to transform a mammalian, insect or microbial host cell. Transformation can be performed using any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus or bacteriophage and transducing a host cell with the construct by transfection procedures known in the art, as exemplified by U.S. Pat. Nos. 4,399,216; 4,912,040; 4,740,461; 4,959,455. The optimal transformation procedure used will depend upon which type of host cell is being transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include, but are not limited to, dextran-mediated trans-

fection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, mixing nucleic acid with positively-charged lipids, and direct microinjection of the DNA into nuclei.

Recombinant expression constructs typically comprise a polynucleotide encoding a polypeptide. The polypeptide may comprise one or more of the following: one or more CDRs such as provided herein; a light chain variable region; a heavy chain variable region; a light chain constant region; a heavy chain constant region (e.g., C_H1 , C_H2 and/or C_H3); and/or another scaffold portion of an IL-23 antigen binding protein. These nucleic acid sequences are inserted into an appropriate expression vector using standard ligation techniques. In one embodiment, the heavy or light chain constant region is appended to the C-terminus of a heavy or light chain variable region provided herein and is ligated into an expression vector. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery, permitting amplification and/or expression of the gene can occur). In some embodiments, vectors are used that employ protein-fragment complementation assays using protein reporters, such as dihydrofolate reductase (see, for example, U.S. Pat. No. 6,270,964). Suitable expression vectors can be purchased, for example, from Invitrogen Life Technologies (Carlsbad, Calif.) or BD Biosciences (San Jose, Calif.). Other useful vectors for cloning and expressing the antibodies and fragments include those described in Bianchi and McGrew, 2003, *Biotech. Biotechnol. Bioeng.* 84:439-44. Additional suitable expression vectors are discussed, for example, in *Methods Enzymol.*, vol. 185 (D. V. Goeddel, ed.), 1990, New York: Academic Press.

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as "flanking sequences" in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the polynucleotide encoding the polypeptide to be expressed, and a selectable marker element. The expression vectors that are provided may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Optionally, the vector may contain a "tag"-encoding sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the IL-23 antigen binding protein coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another "tag" such as FLAG®, HA (hemagglutinin influenza virus), or myc, for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification or detection of the IL-23 antigen binding protein from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an

affinity matrix. Optionally, the tag can subsequently be removed from the purified IL-23 antigen binding protein by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), synthetic or native. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

Flanking sequences useful in the vectors may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Whether all or only a portion of the flanking sequence is known, it may be obtained using polymerase chain reaction (PCR) and/or by screening a genomic library with a suitable probe such as an oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Qiagen, Chatsworth, Calif.), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, Mass.) is suitable for most gram-negative bacteria, and various viral origins (e.g., SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it also contains the virus early promoter).

A transcription termination sequence is typically located 3' to the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for pro-

karyotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex or defined media. Specific selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. Advantageously, a neomycin resistance gene may also be used for selection in both prokaryotic and eukaryotic host cells.

Other selectable genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are required for production of a protein critical for growth or cell survival are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and promoterless thymidine kinase genes. Mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selectable gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively increased, thereby leading to the amplification of both the selectable gene and the DNA that encodes another gene, such as an antigen binding protein that binds to IL-23. As a result, increased quantities of a polypeptide such as an antigen binding protein are synthesized from the amplified DNA.

A ribosome-binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the polypeptide to be expressed.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various pre- or pro-sequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add prosequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein), one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired polypeptide, if the enzyme cuts at such area within the mature polypeptide.

Expression and cloning will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding an IL-23 antigen binding protein. Promoters are untranscribed sequences located upstream (i.e., 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, uniformly transcribe a gene to which they are operably linked, that is, with little or no control over gene expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding a heavy chain variable region or a light chain variable region of an IL-23 antigen binding protein by

removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector.

Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus, and Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

Additional promoters which may be of interest include, but are not limited to: SV40 early promoter (Benoist and Chambon, 1981, *Nature* 290:304-310); CMV promoter (Thomsen et al., 1984, *Proc. Natl. Acad. U.S.A.* 81:659-663); the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, *Cell* 22:787-797); herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-1445); promoter and regulatory sequences from the metallothioneine gene (Prinster et al., 1982, *Nature* 296:39-42); and prokaryotic promoters such as the beta-lactamase promoter (Villakamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731); or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region that is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region that is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122); the immunoglobulin gene control region that is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444); the mouse mammary tumor virus control region that is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495); the albumin gene control region that is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276); the alpha-feto-protein gene control region that is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 253:53-58); the alpha 1-antitrypsin gene control region that is active in liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171); the beta-globin gene control region that is active in myeloid cells (Mogam et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94); the myelin basic protein gene control region that is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); the myosin light chain-2 gene control region that is active in skeletal muscle (Sani, 1985, *Nature* 314:283-286); and the gonadotropic releasing hormone gene control region that is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378).

An enhancer sequence may be inserted into the vector to increase transcription by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent, having been found at positions both 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, how-

ever, an enhancer from a virus is used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers known in the art are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be positioned in the vector either 5' or 3' to a coding sequence, it is typically located at a site 5' from the promoter. A sequence encoding an appropriate native or heterologous signal sequence (leader sequence or signal peptide) can be incorporated into an expression vector, to promote extracellular secretion of the antibody. The choice of signal peptide or leader depends on the type of host cells in which the antibody is to be produced, and a heterologous signal sequence can replace the native signal sequence. Examples of signal peptides that are functional in mammalian host cells include the following: the signal sequence for interleukin-7 described in U.S. Pat. No. 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., 1984, *Nature* 312:768; the interleukin-4 receptor signal peptide described in EP Patent No. 0367 566; the type I interleukin-1 receptor signal peptide described in U.S. Pat. No. 4,968,607; the type II interleukin-1 receptor signal peptide described in EP Patent No. 0 460 846.

After the vector has been constructed, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an antigen binding protein into a selected host cell may be accomplished by well known methods including transfection, infection, calcium phosphate co-precipitation, electroporation, microinjection, lipofection, DEAE-dextran mediated transfection, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001).

A host cell, when cultured under appropriate conditions, synthesizes protein that can be subsequently collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and constitutively produce antigen binding proteins with IL-23 binding properties. In another embodiment, a cell line from the B cell lineage that does not make its own antibody but has a capacity to make and secrete a heterologous antibody can be also selected.

Use of Human IL-23 Antigen Binding Proteins for Diagnostic and Therapeutic Purposes

Antigen binding proteins are useful for detecting IL-23 in biological samples and identification of cells or tissues that produce IL-23. Antigen binding proteins that specifically

bind to IL-23 may be used in diagnosis and/or treatment of diseases related to IL-23 in a patient in need thereof. For one, the IL-23 antigen binding proteins can be used in diagnostic assays, e.g., binding assays to detect and/or quantify IL-23 expressed in blood, serum, cells or tissue. In addition, IL-23 antigen binding proteins can be used to reduce, inhibit, interfere with or modulate one or more biological activities of IL-23 in a cell or tissue. Thus antigen binding proteins that bind to IL-23 may have therapeutic use in ameliorating diseases related to IL-23.

Indications

The present invention also relates to the use of IL-23 antigen binding proteins for use in the prevention or therapeutic treatment of medical disorders, such as those disclosed herein. The IL-23 antigen binding proteins are useful to treat a variety of conditions in which IL-23 is associated with or plays a role in contributing to the underlying disease or disorder or otherwise contributes to a negative symptom.

Conditions effectively treated by IL-23 antigen binding proteins play a role in the inflammatory response. Such inflammatory disorders include periodontal disease; lung disorders such as asthma; skin disorders such as psoriasis, atopic dermatitis, contact dermatitis; rheumatic disorders such as rheumatoid arthritis, progressive systemic sclerosis (scleroderma); systemic lupus erythematosus; spondyloarthritis including ankylosing spondylitis, psoriatic arthritis, enteropathic arthritis and reactive arthritis. Also contemplated is uveitis including Vogt-Koyanagi-Harada disease, idiopathic anterior and posterior uveitis, and uveitis associated with spondyloarthritis. Use of IL-23 antigen binding proteins is also contemplated for the treatment of autoimmune disorders including multiple sclerosis; autoimmune myocarditis; type 1 diabetes and autoimmune thyroiditis.

Degenerative conditions of the gastrointestinal system are treatable or preventable with IL-23 antigen binding proteins. Such gastrointestinal disorders including inflammatory bowel disease: Crohn's disease, ulcerative colitis and Celiac disease.

Also included are use of IL-23 antigen binding proteins in treatments for graft-versus-host disease, and complications such as graft rejection, resulting from solid organ transplantation, such as heart, liver, skin, kidney, lung or other transplants, including bone marrow transplants.

Also provided herein are methods for using IL-23 antigen binding proteins to treat various oncologic disorders including various forms of cancer including colon, stomach, prostate, renal cell, cervical and ovarian cancers, and lung cancer (SCLC and NSCLC). Also included are solid tumors, including sarcoma, osteosarcoma, and carcinoma, such as adenocarcinoma and squamous cell carcinoma, esophageal cancer, gastric cancer, gall bladder carcinoma, leukemia, including acute myelogenous leukemia, chronic myelogenous leukemia, myeloid leukemia, chronic or acute lymphoblastic leukemia and hairy cell leukemia, and multiple myeloma.

Diagnostic Methods

The antigen binding proteins of the described can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or conditions associated with IL-23. Examples of methods useful in the detection of the presence of IL-23 include immunoassays, such as the enzyme linked immunosorbant assay (ELISA) and the radioimmunoassay (RIA).

For diagnostic applications, the antigen binding protein typically will be labeled with a detectable labeling group. Suitable labeling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{15}N , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I), fluorescent groups (e.g.,

FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the labelling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used.

Other diagnostic methods are provided for identifying a cell or cells that express IL-23. In a specific embodiment, the antigen binding protein is labeled with a labeling group and the binding of the labeled antigen binding protein to IL-23 is detected. In a further specific embodiment, the binding of the antigen binding protein to IL-23 is detected in vivo. In a further specific embodiment, the IL-23 antigen binding protein is isolated and measured using techniques known in the art. See, for example, Harlow and Lane, 1988, *Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor (ed. 1991 and periodic supplements); John E. Coligan, ed., 1993, *Current Protocols In Immunology* New York: John Wiley & Sons.

Other methods provide for detecting the presence of a test molecule that competes for binding to IL-23 with the antigen binding proteins provided. An example of one such assay would involve detecting the amount of free antigen binding protein in a solution containing an amount of IL-23 in the presence or absence of the test molecule. An increase in the amount of free antigen binding protein (i.e., the antigen binding protein not bound to IL-23) would indicate that the test molecule is capable of competing for IL-23 binding with the antigen binding protein. In one embodiment, the antigen binding protein is labeled with a labeling group. Alternatively, the test molecule is labeled and the amount of free test molecule is monitored in the presence and absence of an antigen binding protein.

Methods of Treatment: Pharmaceutical Formulations, Routes of Administration

Pharmaceutical compositions that comprise a therapeutically effective amount of one or a plurality of the antigen binding proteins and a pharmaceutically acceptable excipient, diluent, carrier, solubilizer, emulsifier, preservative, and/or adjuvant are provided. In addition, methods of treating a patient by administering such pharmaceutical composition are included. The term "patient" includes human patients. The terms "treat" and "treatment" encompass alleviation or prevention of at least one symptom or other aspect of a disorder, or reduction of disease severity, and the like. The term "therapeutically effective amount" or "effective amount" refers to the amount of an IL-23 antigen binding protein determined to produce any therapeutic response in a mammal. Such therapeutically effective amounts are readily ascertained by one of ordinary skill in the art.

An antigen binding protein need not affect a complete cure, or eradicate every symptom or manifestation of a disease, to constitute a viable therapeutic agent. As is recognized in the pertinent field, drugs employed as therapeutic agents may reduce the severity of a given disease state, but need not abolish every manifestation of the disease to be regarded as useful therapeutic agents. Similarly, a prophylactically administered treatment need not be completely effective in preventing the onset of a condition in order to constitute a viable prophylactic agent. Simply reducing the impact of a disease (for example, by reducing the number or severity of its symptoms, or by increasing the

effectiveness of another treatment, or by producing another beneficial effect), or reducing the likelihood that the disease will occur or worsen in a subject, is sufficient. Certain methods provided herein comprise administering to a patient an IL-23 antagonist (such as the antigen binding proteins disclosed herein) in an amount and for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of the particular disorder.

As is understood in the pertinent field, pharmaceutical compositions comprising the molecules of the invention are administered to a patient in a manner appropriate to the indication. Pharmaceutical compositions may be administered by any suitable technique, including but not limited to, parenterally, topically, or by inhalation. If injected, the pharmaceutical composition can be administered, for example, via intra-articular, intravenous, intramuscular, intraslesional, intraperitoneal or subcutaneous routes, by bolus injection, or continuous infusion. Localized administration, e.g. at a site of disease or injury is contemplated, as are transdermal delivery and sustained release from implants. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation of the antagonist in aerosol form, and the like. Other alternatives include eyedrops; oral preparations including pills, syrups, lozenges or chewing gum; and topical preparations such as lotions, gels, sprays, and ointments.

Use of antigen binding proteins in ex vivo procedures also is contemplated. For example, a patient's blood or other bodily fluid may be contacted with an antigen binding protein that binds IL-23 ex vivo. The antigen binding protein may be bound to a suitable insoluble matrix or solid support material.

Advantageously, antigen binding proteins are administered in the form of a composition comprising one or more additional components such as a physiologically acceptable carrier, excipient or diluent. Optionally, the composition additionally comprises one or more physiologically active agents for combination therapy. A pharmaceutical composition may comprise an IL-23 antigen binding protein together with one or more substances selected from the group consisting of a buffer, an antioxidant such as ascorbic acid, a low molecular weight polypeptide (such as those having fewer than 10 amino acids), a protein, an amino acid, a carbohydrate such as glucose, sucrose or dextrans, a chelating agent such as EDTA, glutathione, a stabilizer, and an excipient. Neutral buffered saline or saline mixed with conspecific serum albumin are examples of appropriate diluents. In accordance with appropriate industry standards, preservatives such as benzyl alcohol may also be added. The composition may be formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Suitable components are nontoxic to recipients at the dosages and concentrations employed. Further examples of components that may be employed in pharmaceutical formulations are presented in any Remington's Pharmaceutical Sciences including the 21st Ed. (2005), Mack Publishing Company, Easton, Pa.

Kits for use by medical practitioners include an IL-23 antigen binding protein and a label or other instructions for use in treating any of the conditions discussed herein. In one embodiment, the kit includes a sterile preparation of one or more IL-23 binding antigen binding proteins, which may be in the form of a composition as disclosed above, and may be in one or more vials.

Dosages and the frequency of administration may vary according to such factors as the route of administration, the particular antigen binding proteins employed, the nature and

severity of the disease to be treated, whether the condition is acute or chronic, and the size and general condition of the subject. Appropriate dosages can be determined by procedures known in the pertinent art, e.g. in clinical trials that may involve dose escalation studies.

A typical dosage may range from about 0.1 µg/kg to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 0.1 µg/kg up to about 30 mg/kg, optionally from 1 µg/kg up to about 30 mg/kg, optionally from 10 µg/kg up to about 10 mg/kg, optionally from about 0.1 mg/kg to 5 mg/kg, or optionally from about 0.3 mg/kg to 3 mg/kg.

Dosing frequency will depend upon the pharmacokinetic parameters of the particular human IL-23 antigen binding protein in the formulation used. Typically, a clinician administers the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Appropriate dosages may be ascertained through use of appropriate dose-response data. An IL-23 antigen binding protein of the invention may be administered, for example, once or more than once, e.g., at regular intervals over a period of time. In particular embodiments, an IL-23 antigen binding protein is administered over a period of at least a month or more, e.g., for one, two, or three months or even indefinitely. For treating chronic conditions, long-term treatment is generally most effective. However, for treating acute conditions, administration for shorter periods, e.g. from one to six weeks, may be sufficient. In general, the antigen binding protein is administered until the patient manifests a medically relevant degree of improvement over baseline for the chosen indicator or indicators.

It is contemplated that an IL-23 antigen binding protein be administered to the patient in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the patient's illness, disease or condition may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. In one embodiment, an improvement is considered to be sustained if the subject exhibits the improvement on at least two occasions separated by two to four weeks. The degree of improvement generally is determined by a physician, who may make this determination based on signs, symptoms, biopsies, or other test results, and who may also employ questionnaires that are administered to the subject, such as quality-of-life questionnaires developed for a given disease.

Particular embodiments of methods and compositions of the invention involve the use of an IL-23 antigen binding protein and one or more additional IL-23 antagonists, for example, two or more antigen binding proteins of the invention, or an antigen binding protein of the invention and one or more other IL-23 antagonists. Also provided are IL-23 antigen binding proteins administered alone or in combination with other agents useful for treating the condition with which the patient is afflicted. Examples of such agents include both proteinaceous and non-proteinaceous drugs. Such agents include therapeutic moieties having anti-inflammatory properties (for example, non-steroidal anti-inflammatory agents, steroids, immunomodulators and/or other cytokine inhibitors such as those that antagonize, for

example, IFN- γ , GM-CSF, IL-6, IL-8, IL-17, IL-22 and TNFs), or of an IL-23 antigen binding protein and one or more other treatments (e.g., surgery, ultrasound, or treatment effective to reduce inflammation). When multiple therapeutics are co-administered, dosages may be adjusted accordingly, as is recognized or known in the pertinent art. Useful agents that may be combined with IL-23 antigen binding proteins include those used to treat, for example, Crohn's disease or ulcerative colitis, such as aminosalicylate (for example, mesalamine), corticosteroids (including prednisone), antibiotics such as metronidazole or ciprofloxacin (or other antibiotics useful for treating, for example, patients afflicted with fistulas), and immunosuppressives such as azathioprine, 6-mercaptopurine, methotrexate, tacrolimus and cyclosporine. Such agent(s) may be administered orally or by another route, for example via suppository or enema. Agents which may be combined with IL-23 binding proteins in treatment of psoriasis include corticosteroids, calcipotriene and other vitamin D derivatives, acetrelin and other retinoic acid derivatives, methotrexate, tacrolimus, and cyclosporine used topically or systemically. Such agents can be administered simultaneously, consecutively, alternately, or according to any other regimen that allows the total course of therapy to be effective.

In addition to human patients, IL-23 antigen binding proteins are useful in the treatment of non-human animals, such as domestic pets (dogs, cats, birds, primates, etc.), domestic farm animals (horses, cattle, sheep, pigs, birds, etc). In such instances, an appropriate dose may be determined according to the animal's body weight. For example, a dose of 0.2-1 mg/kg may be used. Alternatively, the dose is determined according to the animal's surface area, an exemplary dose ranging from 0.1-20 mg/m², or more preferably, from 5-12 mg/m². For small animals, such as dogs or cats, a suitable dose is 0.4 mg/kg. IL-23 antigen binding protein (preferably constructed from genes derived from the recipient species) is administered by injection or other suitable route one or more times per week until the animal's condition is improved, or it may be administered indefinitely.

The following examples, including the experiments conducted and the results achieved, are provided for illustrative purposes only and are not to be construed as limiting the scope of the appended claims.

EXAMPLES

Example 1

Generation of Human IL-23 Antibodies

XenoMouse™ technology (Amgen, Thousand Oaks, Calif.) was used to develop human monoclonal antibodies that recognize and inhibit native human IL-23 activity while sparing human IL-12. The antibodies also recognize and inhibit recombinant cynomolgous IL-23 but do not recognize murine or rat IL-23.

Antibodies were selected for recognition and complete inhibition of native human IL-23 obtained from human monocyte-derived dendritic cells (MoDCs), using the STAT-luciferase reporter assay described below. Human monocytes were isolated from peripheral blood mononuclear cells from healthy donors using negative selection (Monocyte Isolation Kit II, Miltenyi Biotec, Auburn, Calif.). MoDCs were generated by culturing monocytes with human GM-CSF (50 ng/ml) and human IL-4 (100 ng/ml) for 7 days in RPMI 1640 with 10% fetal bovine serum complete medium. MoDCs were then washed twice with PBS followed by

stimulation with human CD40L (1 μ g/ml) for an additional 48 hours. CD40L-stimulated MoDC supernatant contains IL-23, IL-12 and IL-12/23p40. ELISAs are used to determine the amount of IL-12p70 (R&D System, Minneapolis, Minn.), IL-23 (eBiosciences, San Diego, Calif.) and IL-12/23p40 (R&D Systems). The STAT-luciferase assay responds to IL-23 and not to IL-12 or to free IL-12/23p40, therefore the assay could be used with crude supernatants to assess IL-23 activity. For use in the NK cell assay, described below, the native human IL-23 crude supernatant was purified using an IL-23 affinity column followed by size exclusion chromatography. Concentration was determined using an IL-23 specific ELISA (eBiosciences).

The purified antibody supernatants were also tested against recombinant human (rhu) IL-23 and recombinant cynomolgous (cyno) IL-23 in the STAT-luciferase assay. Of the antibodies tested that completely inhibited recombinant human IL-23, only half of those antibodies recognized and completely inhibited native human IL-23. Recognition and complete inhibition of recombinant human IL-23 was not predictive of, nor correlated to, recognition and complete inhibition of native human IL-23. As shown in FIGS. 1A and 1B, of the antibody supernatants that completely inhibited recombinant human IL-23, only half of those antibodies completely inhibited native human IL-23. Those antibodies that recognized and completely inhibited native human IL-23 were selected for further characterization.

Example 2

Functional Assays

a) STAT-Luciferase Assay

It is known that IL-23 binds its heterodimeric receptor and signals through JAK2 and Tyk2 to activate STAT 1, 3, 4 and 5. In this assay, cells transfected with a STAT/luciferase reporter gene are used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced bioactivity.

Chinese hamster ovary cells expressing human IL-23 receptor are transiently transfected with STAT-luciferase reporter overnight. IL-23 antibodies are serially diluted (12 points of 1:4 serial dilutions starting at 37.5 μ g/ml) into 96 well plates. Native human IL-23 (preparation method is described in Example 1) is added to each well at a concentration of 2 ng/ml and incubated at room temperature for 15-20 minutes. The transiently transfected cells are added (8×10^3 cells) to a final volume of 100 μ L/well and incubated for 5 hours at 37° C., 10% CO₂. Following incubation, cells are lysed using 100 μ L/well Glo Lysis buffer (1 $\times$) (Promega, Madison, Wis.) at room temperature for 5 minutes. Fifty microliters of cell lysate is added to a 96 well plate along with 50 μ L Bright-Glo luciferase substrate (Promega) and read on a luminometer.

Statistical analysis can be performed using GraphPad PRISM software (GraphPad Software, La Jolla, Calif.). Results can be expressed as the mean $\pm$ standard deviation (SD).

As seen in TABLE 5, all IL-23 antibodies potently and completely inhibited native human IL-23-induced STAT/luciferase reporter in a dose dependent manner. The antibodies also potently and completely inhibited recombinant human (rhu) IL-23 and recombinant cyno (cyno) IL-23. The antibodies all had IC₅₀ values in the picomolar range.

TABLE 5

Table of mean IC ₅₀ (pM) values for IL-23 antibodies in the STAT-luciferase assay.						
antibody	Native huIL-23		rhuIL-23		Cyno IL-23	
	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats
A	114 +/- 70	3	190 +/- 99	3	379 +/- 213	3
B	45 +/- 5	4	100 +/- 59	4	130 +/- 60	3
C	107 +/- 31	3	211 +/- 93	3	376 +/- 89	3
D	65 +/- 5	3	107 +/- 30	3	184 +/- 77	3
E	140 +/- 52	3	142 +/- 52	3	188 +/- 59	3
F	86 +/- 47	4	187 +/- 116	4	366 +/- 219	4
G	156 +/- 74	5	296 +/- 133	5	421 +/- 174	5
H	192 +/- 35	4	253 +/- 184	4	1024 +/- 533	4
I	208 +/- 33	3	338 +/- 140	3	650 +/- 42	3
J	83 +/- 54	2	36 +/- 6	2	56 +/- 2	2
K	71 +/- 38	3	43 +/- 20	3	61 +/- 10	3
L	113 +/- 80	3	23 +/- 7	3	47 +/- 1	3
M	34 +/- 11	2	40 +/- 8	2	56 +/- 6	2
N	361 +/- 164	3	145	1	238	1

20

b) NK Cell Assay

It is known that IL-23 acts on natural killer cells to induce expression of pro-inflammatory cytokines, such as interferon γ (IFN γ). In this assay, human primary natural killer (NK) cells are used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced IFN γ activity in cells expressing the native receptor for human IL-23.

NK cells are isolated from multiple human donors via negative selection (NK Cell Isolation Kit, Miltenyi Biotec, Auburn, Calif.). Purified NK cells (1×10^6 cells/ml) are added to 6 well plates in RPMI 1640 plus 10% fetal bovine serum complete medium supplemented with recombinant human IL-2 (10 ng/ml, R&D Systems, Minneapolis, Minn.),

cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range. The assay was performed on a subset of antibodies using native human IL-23 (30 μ g/ml, preparation method is described in Example 1) and rhuIL-18 (40 ng/ml, R&D Systems) and yielded the results shown in TABLE 6. Consistent with the selection for IL-23 specific antibodies, these anti-IL-23 antibodies had no effect on IL-12 stimulated IFN γ production in NK cells using the assay described above, whereas an IL-12p35 specific neutralizing antibody, mAb219 (R&D Systems, Minneapolis, Minn.) potentially inhibited recombinant human IL-12.

TABLE 6

Table of mean IC ₅₀ (pM) values for IL-23 antibodies in the NK cell assay.						
antibody	Native huIL-23		rhuIL-23		Cyno IL-23	
	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats
A			42 +/- 12	2	31 +/- 21	2
B	85 +/- 30	2	48 +/- 30	3	19 +/- 8	2
C			32 +/- 19	4	29 +/- 16	2
D			37 +/- 21	2	29 +/- 19	2
E	158 +/- 50	2	57 +/- 14	3	21 +/- 3	2
F			25 +/- 15	2	21 +/- 17	2
G	152 +/- 72	2	45 +/- 30	3	23 +/- 8	2
H			29 +/- 28	2	33 +/- 17	2
I			69	1	52	1
J			4 +/- 3	2	5 +/- 3	2
K			7 +/- 2	2	8 +/- 6	2
L			3 +/- 1	2	4 +/- 1	2
M			8	1	12	1

to a final volume of 10 ml/well. Cells are cultured for 7 days at 37° C., 5% CO₂. The IL-2-activated NK cells are then stimulated with rhuIL-23 or cyno IL-23 (10 ng/ml) and recombinant human IL-18 (20 ng/ml, R&D Systems, Minneapolis, Minn.) in the presence of serial dilutions (11 points of 1:3 serial dilutions starting at 3 μ g/ml) of IL-23 antibodies for 24 hours. IFN γ levels are measured in the supernatant by IFN γ ELISA (R&D Systems, Minneapolis, Minn.) according to manufacturer's instructions.

Statistical analysis can be performed using GraphPad PRISM software. Results can be expressed as the mean $\pm$ standard deviation (SD).

As seen in TABLE 6, all antibodies potentially inhibited rhuIL-23 and cyno IL-23-induced IFN γ expression in NK

c) Human Whole Blood Assay

Human whole blood is collected from multiple healthy donors using Refludan® (Bayer Pittsburgh, Pa.) as an anti-coagulant. The final concentration of Refludan® in whole blood is 10 μ g/ml. A stimulation mixture of rhuIL-23 or cyno IL-23 (final concentration 1 ng/ml)+rhuIL-18 (final concentration 20 ng/ml)+rhuIL-2 (final concentration 5 ng/ml) in RPMI 1640+10% FBS, is added to a 96 well plate, final volume 20 μ l/well. Serially diluted IL-23 antibodies (11 points of 1:3 serial dilutions starting from 3 μ g/ml) are added at 20 μ l/well and incubated with the stimulation mixture for 30 minutes at room temperature. Whole blood is then added (120 μ l/well) and the final volume adjusted to 200 μ l/well

61

with RPMI 1640+10% FBS. The final concentration of whole blood is 60%. The plates are incubated for 24 hours at 37° C., 5% CO₂. Cell free supernatants are harvested and IFN γ levels are measured from the supernatants by IFN γ ELISA (R&D Systems) according to manufacturer's instructions.

Statistical analysis can be performed using GraphPad PRISM software. Results can be expressed as the mean $\pm$ standard deviation (SD).

As seen in TABLE 7, all antibodies potentially inhibited rhuIL-23-induced and cyno-IL-23-induced IFN γ expression in whole blood cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range.

TABLE 7

Table of mean IC ₅₀ (pM) values for IL-23 antibodies in the IFN γ human whole blood assay				
antibody	rhuIL-23		Cyno IL-23	
	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats
B	117 +/- 94	7	161 +/- 95	6
E	29 +/- 8	3	54 +/- 33	3
G	53 +/- 13	3	93 +/- 44	3
F	66 +/- 13	3	166 +/- 189	3
D	88 +/- 6	3	110 +/- 14	3
C	97 +/- 31	3	186 +/- 194	3

d) IL-22 Assay

It is known that IL-23 is a potent inducer of proinflammatory cytokines. IL-23 acts on activated and memory T cells and promotes the survival and expansion of Th17 cells which produce proinflammatory cytokines including IL-22. In this assay, human whole blood is used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced IL-22 production.

A whole blood assay is conducted in the same manner as described above with the modification of using rhuIL-23 or cynoIL-23 at 1 ng/ml and rhuIL-18 at 10 ng/ml to induce IL-22 production. IL-22 concentration is determined by IL-22 ELISA (R&D Systems, Minneapolis, Minn.).

As seen in TABLE 8, the antibodies potentially inhibited rhuIL-23-induced and cyno IL-23-induced IL-22 production in whole blood cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range.

TABLE 8

Table of mean IC ₅₀ (pM) values for IL-23 antibodies in the IL-22 human whole blood assay				
antibody	rhuIL-23		Cyno IL-23	
	IC ₅₀ +/- SD	Repeats	IC ₅₀ +/- SD	Repeats
B	117 +/- 68	4	113 +/- 65	3
E	87 +/- 109	3	56 +/- 60	3
G	83 +/- 59	3	66 +/- 45	3

Example 3

Determining the Equilibrium Dissociation Constant (K_D) for Anti-IL-23 Antibodies Using KinExA Technology

Binding affinity of rhuIL-23 to IL-23 antibodies is evaluated using a kinetic exclusion assay (KinExA assay, Sapi-dyne Instruments, Inc., Boise, Id.). Normal human serum

62

(NHS)-activated Sepharose 4 fast flow beads (Amersham Biosciences, part of GE Healthcare, Uppsala, Sweden), are pre-coated with rhuIL-23 and blocked with 1 M Tris buffer with 10 mg/mL BSA. 50 pM of IL-23 antibody is incubated with rhuIL-23 (12 points of 1:2 dilutions starting from 800 pM) at room temperature for 72 hours before it is run through the rhuIL-23-coated Sepharose beads. The amount of the bead-bound antibody was quantified by fluorescent (Cy5) labeled goat anti-human-Fc antibody (Jackson Immuno Research, West Grove, Pa.). The binding signal is proportional to the amount of free antibody at equilibrium.

The dissociation equilibrium constant (K_D) and the association rate (K_{on}) are obtained from curve fitting using KinExA Pro software. The dissociation rate (K_{off}) is derived from: K_D=K_{off}/K_{on}.

As seen in TABLE 9, the antibodies have high affinity for binding to human IL-23. All had K_D values in the low to sub pM range.

TABLE 9

Table of K _D (pM), K _{on} (1/MS) and K _{off} (1/s) rates			
Antibody	KD (pM)	Kon (1/MS)	Koff (1/s)
E	0.131	9.12E+05	1.4E-07
D	0.126	1.72E+06	2.2E-07
B	3.99	1.17E+06	4.7E-06
C	2.56	1.36E+06	4.1E-06
F	2.62	5.69E+05	1.5E-06
L	1.08	3.34E+06	3.7E-06
G	2.00	4.00E+05	8.1E-07

Example 4

Structure Determination Using X-Ray Crystallography

One way to determine the structure of an antibody-antigen complex is by using X-ray crystallography, see for example, Harlow and Lane Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), p. 23. The crystal structure of IL-23 has been determined, (see Lupardus and Garcia, J Mol Biol, 2008, 382: 931-941) and the crystal structure of an IL-23/Fab complex has been disclosed, (see Beyer et al. J Mol Biol, 2008, 382(4): 942-55). Structural determination of IL-23 with Fab fragments of antibodies claimed herein was obtained using X-ray crystallography.

Protein for Crystallization

A recombinantly derived human IL-23 heterodimer was used for the crystallization studies (see Beyer et al., supra). The sequence of the human p19 subunit comprised of residues 20-189 of SEQ ID NO: 145, the signal sequence of SEQ ID NO:154 and a C-terminal 6-His tag SEQ ID NO:155. The sequence of the human p40 subunit was mutated from asparagine to glutamine at position 222 of SEQ ID NO:147 in order to prevent glycosylation at this site (Beyer, et al., supra).

Fabs derived from Antibody B and Antibody E were expressed on an IgG1 scaffold that incorporated a caspase cleavage site. The Fabs were processed by means of protease cleavage.

Complex Formation and Crystallization

The IL-23-Antibody B Fab complex was made by mixing a 2 $\times$ molar excess of the Antibody B Fab with the human heterodimeric IL-23 described above. The complex was purified by size exclusion chromatography to remove excess

Antibody B Fab and concentrated to ~12 mg/ml for crystallization. The IL-23-Antibody B Fab complex crystallized in 0.1 M Hepes pH 7, 8% PEG 8000.

The IL-23-Antibody E Fab complex was made by mixing a 2× molar excess of the Antibody E Fab with the human heterodimeric IL-23 described above. The complex was methylated using a JBS Methylation Kit according to manufacturer's instructions (Jena Bioscience, Jena, Germany). The complex was then treated with PNGase to deglycosylate the protein. Following these treatments, the complex was purified by size exclusion chromatography to remove excess Antibody E Fab and concentrated to 13.5 mg/ml for crystallization. The IL-23-Antibody E Fab complex crystallized in 0.1 M Tris pH 8.5, 0.2 M magnesium chloride, 15% PEG 4000.

Data Collection and Structure Determination

IL-23-Antibody B Fab crystals grew in the $P2_1$ space group with unit cell dimensions $a=70.93$, $b=71.27$, $c=107.37$ Å, $\beta=104.98^\circ$ and diffract to 2.0 Å resolution. The IL-23-Antibody B Fab structure was solved by molecular replacement with the program MOLREP (CCP4, The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr*, 1994, 50(Pt 5): p. 760-3) using the IL-23 structure (Beyer et al. *supra*) as the starting search model. Keeping the IL-23 solution fixed, an antibody variable domain was used as a search model. Keeping the IL-23-antibody variable domain solution fixed, an antibody constant domain was used as a search model. The complete structure was improved with multiple rounds of model building with Quanta and refinement with *cnx* (Brunger, et al., *Acta Crystallogr D Biol Crystallogr*, 1998, 54(Pt 5): p. 905-21).

Distances between protein atoms were calculated using the program PyMOL (DeLano, W. L. The PyMOL Graphics System. Palo Alto, 2002) (Schrodinger, LLC; New York, N.Y.). Amino acids were chosen if at least one atom was located within the required distance threshold to the partner protein.

Boundaries of the A, B, C and D helices of the p19 subunit of IL-23 when bound to the Antibody B Fab include A helix residues 28-47, B helix residues 86-105, C helix residues 119-134 and D helix residues 154-187 of SEQ ID NO:145.

The regions of interaction on the IL-23p19 subunit when bound to the Antibody B Fab include residues within Ser46-Glu58, Glu112-Glu123 and Pro155-Phe163 of SEQ ID NO:145.

IL-23p19 subunit amino acid residues with atoms 4 Å or less from the Antibody B Fab include Ser46, Ala47, His48, Pro49, Leu50, His53, Met54, Asp55, Glu58, Pro113, Ser114, Leu115, Leu116, Pro120, Val121, Trp156, Leu159, Leu160, Arg162 and Phe163 of SEQ ID NO:145. IL-23p19 amino acid residues with atoms between 4 Å and 5 Å from the Antibody B Fab include Val51, Arg57, Glu112, Asp118, Ser119, Gln123, Pro155 of SEQ ID NO:145.

IL-23p40 subunit amino acid residues with atoms 4 Å or less from the Antibody B Fab include Glu 122 and Lys 124 of SEQ ID NO:147.

The Antibody B Fab heavy chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Gly32, Gly33, Tyr34, Tyr35, His54, Asn58, Thr59, Tyr60, Lys66, Arg101, Gly102, Phe103, Tyr104 and Tyr105 of SEQ ID NO:46. The Antibody B Fab heavy chain amino acid residues with atoms ≤5 Å from the IL-23 heterodimer include Ser31, Gly32, Gly33, Tyr34, Tyr35, His54, Ser56, Asn58, Thr59, Tyr60, Lys66, Arg101, Gly102, Phe103, Tyr104 and Tyr105 of SEQ ID NO:46.

The Antibody B Fab light chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Ser30, Ser31, Trp32, Tyr49, Ser52, Ser53, Ala91, Asn92, Ser93, Phe94, and Phe96 of SEQ ID NO:15. The Antibody B Fab light chain amino acid residues with atoms ≤5 Å from the IL-23 heterodimer include Ser30, Ser31, Trp32, Tyr49, Ala50, Ser52, Ser53, Ser56, Ala91, Asn92, Ser93, Phe94, and Phe96 of SEQ ID NO:15.

The IL-23-Antibody E Fab complex crystals grew in the $P222_1$ space group with unit cell dimensions $a=61.60$, $b=97.59$, $c=223.95$ Å and diffract to 3.5 Å resolution. The IL-23-Antibody E Fab complex structure was solved by molecular replacement with the program Phaser (CCP4, *supra*) using the IL-23 structure, an antibody variable domain, and an antibody constant domain as the three starting search models, as described above. The complete structure was improved with multiple rounds of model building with Quanta and refinement with *cnx* (Brunger, et al., *supra*). The Antibody E Fab constant domain was left out of the final refined structure due to very poor electron density for that portion of the protein.

The regions of interaction on the IL-23p19 subunit identified when bound to the Antibody E Fab include residues within Ser46-His53, Glu112-Val120 and Trp156-Phe163 of SEQ ID NO:145.

IL-23p19 amino acid residues with atoms 4 Å or less from the Antibody E Fab include Ser46, Ala47, His48, Pro49, Leu50, Glu112, Pro113, Ser114, Leu115, Leu116, Pro117, Asp118, Ser119, Pro120, Trp156, Leu159, Leu160 and Phe163 of SEQ ID NO: 145. IL-23p19 amino acid residues with atoms between 4 Å and 5 Å from the Antibody E Fab include His53 of SEQ ID NO:145.

IL-23p40 amino acid residues with atoms 4 Å or less from the Antibody E Fab include Lys121, Glu 122, Pro123 and Asn 125 of SEQ ID NO:147.

The Antibody E Fab heavy chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Gly26, Phe27, Thr28, Ser31, Tyr53, Tyr59, Tyr102, Ser104, Ser105, Trp106, Tyr107, and Pro108 of SEQ ID NO:31. The Antibody E Fab heavy chain amino acid residues with atoms ≤5 Å from the IL-23 heterodimer include Gln1, Gly26, Phe27, Thr28, Ser30, Ser31, Tyr32, Trp52, Tyr53, Tyr59, Arg100, Tyr102, Thr103, Ser104, Ser105, Trp106, Tyr107, and Pro108 of SEQ ID NO:31.

The Antibody E Fab light chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Ala31, Gly32, Tyr33, Asp34, Tyr51, Gly52, Asn55, Lys68, and Tyr93 of SEQ ID NO:1. The Antibody B Fab light chain amino acid residues with atoms ≤5 Å from the IL-23 heterodimer include Thr29, Ala31, Gly32, Tyr33, Asp34, Tyr51, Gly52, Asn55, Lys68, Tyr93, and Trp100 of SEQ ID NO:1.

Example 5

Determination of IL-23-Antibody Complex Contact Residues Through Solvent Accessible Surface Area Differences

The residue contacts in the paratope (the portion of the antibody that recognizes the antigen) and the portion of the antigen that it binds bound by the paratope in a human IL-23-Antibody B Fab complex and in a human IL-23-Antibody E Fab complex were determined using solvent accessible surface area differences. The solvent accessible

surface area calculations were performed using Molecular Operating Environment (Chemical Computing Group, Montreal, Quebec).

The solvent accessible surface area differences of the paratope residues in the IL-23-Antibody B Fab complex were calculated by setting the Antibody B Fab residues as the desired set. The structural information obtained in Example 4 for the IL-23-Antibody B Fab complex was used and the residue solvent accessible surface area of the amino acid residues of the Antibody B Fab in the presence of the IL-23 heterodimer were calculated and represent the “bound areas” for the set.

The residue solvent accessible surface area of each of the Antibody B Fab residues in the absence of the IL-23 antigen were calculated and represent the “free areas” of the set.

The “bound areas” were then subtracted from the “free areas” resulting in the “solvent exposed surface area difference” for each residue in the set. The Antibody B Fab residues that had no change in surface area, or a zero difference, had no contact with the residues of the IL-23 antigen when complexed. The Antibody B Fab residues that had a difference value $\geq 10 \text{ \AA}^2$ were considered to be in significant contact with residues in the IL-23 antigen such that these Antibody B Fab residues were at least partially to completely occluded when the Antibody B Fab was bound to human IL-23. This set of Antibody B Fab residues make up the “covered patch”, the residues involved in the structure of the interface when Antibody B Fab is bound to human IL-23, see Tables 10 and 11. The Antibody B Fab residues in this covered patch may not be involved in binding interactions with residues of the IL-23 antigen, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody B Fab to human IL-23. With the exception of Tyr49, all of the residues are located in the CDR regions of the Antibody B Fab light and heavy chains. These residues were also within 5 Å or less of the IL-23 antigen when bound to the Antibody B Fab, as described in Example 4.

TABLE 10

Solvent Accessibility Surface Area Differences for Antibody B Fab Light Chain		
Residue AHO Number	Residue Position SEQ ID NO: 15	Solvent exposed surface area difference ($\text{\AA}^2$)
Ser32	Ser30	44.9
Ser33	Ser31	41.1
Trp40	Trp32	79.0
Tyr57	Tyr49	40.7
Ala58	Ala50	20.3
Ser68	Ser52	43.6
Ser69	Ser53	38.9
Ser72	Ser56	19.1
Asn110	Asn92	34.0
Phe135	Phe94	51.4

TABLE 11

Solvent Accessibility Surface Area Differences for Antibody B Fab Heavy Chain		
Residue AHO Number	Residue Position SEQ ID NO: 46	Solvent exposed surface area difference ($\text{\AA}^2$)
Ser33	Ser31	18.2
Gly34	Gly32	49.5
Gly38	Gly33	33.8
Tyr39	Tyr34	51.4

TABLE 11-continued

Solvent Accessibility Surface Area Differences for Antibody B Fab Heavy Chain		
Residue AHO Number	Residue Position SEQ ID NO: 46	Solvent exposed surface area difference ($\text{\AA}^2$)
Tyr40	Tyr35	30.7
His59	His54	29.5
Asn67	Asn58	66.7
Thr68	Thr59	26.0
Tyr69	Tyr60	59.4
Lys75	Lys66	32.6
Arg110	Arg101	47.2
Gly111	Gly102	21.7
Phe112	Phe103	35.5
Tyr133	Tyr104	83.0
Tyr134	Tyr105	91.7

The solvent accessible surface area differences of the residues in the IL-23-Antibody E Fab complex were calculated as described above. The Antibody E Fab residues that had a difference value $\geq 10 \text{ \AA}^2$ were considered to be in significant contact with residues in the IL-23 antigen and these Antibody E Fab residues were at least partially to completely occluded when the Antibody E Fab was bound to human IL-23. This set of Antibody E Fab residues make up the covered patch, the residues involved in the structure of the interface when the Antibody E Fab is bound to human IL-23, see Tables 12 and 13. The Antibody E Fab residues in this covered patch may not be involved in binding interactions with residues of the IL-23 antigen, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody E Fab to human IL-23. For the most part, these covered patch residues were located within the CDR regions of the Antibody E Fab heavy and light chains. These residues were also within 5 Å or less of the IL-23 antigen when bound to the Antibody E Fab, as described in Example 4.

TABLE 12

Solvent Accessibility Surface Area Differences for Antibody E Fab Light Chain		
Residue AHO Number	Residue Position SEQ ID NO: 1	Solvent exposed surface area difference ($\text{\AA}^2$)
Ala33	Ala31	11.6
Gly34	Gly32	51.2
Tyr39	Tyr33	47.2
Asp40	Asp34	36.8
Tyr57	Tyr51	16.1
Gly58	Gly52	11.1
Asn69	Asn55	29.4
Lys82	Lys68	20.1
Tyr109	Tyr93	27.3
Ser135	Ser98	11.3

TABLE 13

Solvent Accessibility Surface Area Differences for Antibody E Fab Heavy Chain		
Residue AHO Number	Residue Position SEQ ID NO: 31	Solvent exposed surface area difference ($\text{\AA}^2$)
Gln1	Gln1	41.1
Gly27	Gly26	24.6
Thr30	Thr28	82.2
Ser33	Ser31	40.7
Tyr39	Tyr32	30.7

67

TABLE 13-continued

Solvent Accessibility Surface Area Differences for Antibody E Fab Heavy Chain		
Residue AHO Number	Residue Position SEQ ID NO: 31	Solvent exposed surface area difference (Å ²)
Trp59	Trp52	11.3
Tyr60	Tyr53	44.7
Tyr69	Tyr59	42.4
Lys86	Lys76	17.4
Gly111	Gly101	12.8
Tyr112	Tyr102	103.1
Ser114	Ser104	21.0
Ser115	Ser105	91.4
Trp131	Trp106	145.0
Tyr132	Tyr107	71.6
Pro133	Pro108	20.4

The solvent accessible surface area differences of the portion of the IL-23 heterodimer bound by the paratope of the Antibody B Fab were calculated by setting the IL-23 heterodimer residues as the desired set. The structural information obtained in Example 4 for the Antibody B Fab-IL-23 complex was used and the residue solvent accessible surface area of the amino acid residues of the IL-23 heterodimer in the presence of the Antibody B Fab were calculated and represent the bound areas for the set.

The residue solvent accessible surface area of each of the IL-23 heterodimer residues in the absence of the Antibody B Fab were calculated and represent the free areas of the set.

As described above, the bound areas were subtracted from the free areas resulting in the solvent exposed surface area difference for each IL-23 residue. The IL-23 heterodimer residues that had no change in surface area, or a zero difference, had no contact with the residues of the Antibody B Fab when complexed. The IL-23 heterodimer residues that had a difference value ≥ 10 Å² were considered to be in significant contact with residues of the Antibody B Fab and these 11-23 heterodimer residues were at least partially to completely occluded when the human IL-23 heterodimer was bound to the Antibody B Fab. This set of IL-23 heterodimer residues make up the covered patch, the residues involved in the structure of the interface when the human IL-23 heterodimer is bound to the Antibody E Fab, see Table 14. The 11-23 heterodimer residues in this covered patch may not all be involved in binding interactions with residues on the Antibody B Fab, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody B Fab to human IL-23. These residues are also within 4 Å or less from the Antibody B Fab, as described Example 4.

TABLE 14

Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues	
	Solvent exposed surface area difference (Å ²)
p19 residues (SEQ ID NO: 145)	
Ser46	26.5
Ala47	12.7
Pro49	59.6
Leu50	122.2
His53	47.8
Met54	13.9
Asp55	20.5
Arg57	14.6

68

TABLE 14-continued

Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues	
	Solvent exposed surface area difference (Å ²)
Glu58	96.5
Glu112	29.7
Pro113	64.8
Ser114	30.0
Leu115	31.4
Leu116	60.0
Asp118	14.4
Ser119	19.7
Pro120	64.7
Pro155	19.4
Trp156	61.9
Leu159	72.8
Leu160	27.0
Arg162	14.4
Phe163	67.5
p40 residues (SEQ ID NO: 147)	
Glu122	29.1
Lys124	60.9

The solvent accessible surface area differences of the portion of the IL-23 heterodimer bound by the paratope of the Antibody E Fab were calculated as described above. The IL-23 heterodimer residues that had a difference value ≥ 10 Å² were considered to be in significant contact with residues of the Antibody E Fab and these 11-23 heterodimer residues were at least partially to completely occluded when the human IL-23 heterodimer was bound to the Antibody E Fab. This set of IL-23 heterodimer residues make up the covered patch, the residues involved in the structure of the interface when the human IL-23 heterodimer is bound to the Antibody E Fab, see Table 15. The 11-23 heterodimer residues in this covered patch may not all be involved in binding interactions with residues on the Antibody E Fab, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody E Fab to human IL-23. These residues are also within 5 Å or less from the Antibody E Fab, as described in Example 4.

TABLE 15

Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues	
	Solvent exposed surface area difference (Å ²)
p19 residues (SEQ ID NO: 145)	
Ser46	18.7
Ala47	14.9
Pro49	79.8
Leu50	99.5
His53	61.2
Glu112	62.8
Pro113	45.7
Ser114	69.5
Leu115	50.3
Leu116	127.2
Pro117	54.1
Asp118	37.0
Pro120	18.8
Pro155	16.9
Trp156	140.7
Leu159	21.8
Leu160	17.0
Phe163	56.6

69

TABLE 15-continued

Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues	
	Solvent exposed surface area difference (Å ²)
p40 residues (SEQ ID NO: 147)	
Lys121	86.2
Glu122	21.8

5

70

TABLE 15-continued

Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues	
	Solvent exposed surface area difference (Å ²)
Pro123	22.1
Asn125	26.7
Arg283	22.6

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 155

<210> SEQ ID NO 1

<211> LENGTH: 111

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Thr Gly Ala Gly
20 25 30

Tyr Asp Val His Trp Tyr Gln Gln Val Pro Gly Thr Ala Pro Lys Leu
35 40 45

Leu Ile Tyr Gly Ser Gly Asn Arg Pro Ser Gly Val Pro Asp Arg Phe
50 55 60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu
65 70 75 80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser
85 90 95

Leu Ser Gly Trp Val Phe Gly Gly Gly Thr Arg Leu Thr Val Leu
100 105 110

<210> SEQ ID NO 2

<211> LENGTH: 333

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

cagtctgtgc tgacgcagcc gccctcagtg tctggggccc cagggcagag ggccaccatc 60

tctgcactg ggagcagctc caacaccggg gcagggtatg atgtacactg gtaccagcaa 120

gttcaggaa cagcccccac actcctcatt tatggtagcg gcaatcggcc ctccaggggtc 180

cctgaccgat tctctggctc caagtctggc acctcagcct cctgggcat cactggactc 240

caggctgagg atgaggctga ttattactgc cagtcctatg acagcagcct gagtggttgg 300

gtgttcggcg gagggaccag gctgaccgtc ctg 333

<210> SEQ ID NO 3

<211> LENGTH: 111

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1 5 10 15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly
20 25 30

-continued

<210> SEQ ID NO 7
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

```

Gln Pro Val Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala
1      5      10      15
Ser Val Thr Leu Thr Cys Thr Leu Asn Ser Gly Tyr Ser Asp Tyr Lys
      20      25      30
Val Asp Trp Tyr Gln Gln Arg Pro Gly Lys Gly Pro Arg Phe Val Met
      35      40      45
Arg Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Asp Gly Ile Pro
      50      55      60
Asp Arg Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Tyr Leu Thr Ile
      65      70      75      80
Lys Asn Ile Gln Glu Glu Asp Glu Ser Asp Tyr His Cys Gly Ala Asp
      85      90      95
His Gly Ser Gly Ser Asn Phe Val Tyr Val Phe Gly Thr Gly Thr Lys
      100     105     110
Val Thr Val Leu
      115

```

<210> SEQ ID NO 8
 <211> LENGTH: 348
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```

cagcctgtgc tgactcagcc accttctgca tcagcctccc tgggagcctc ggtcacactc      60
acctgcaccc tgaacagcgg ctacagtgat tataaagtgg actggtacca gcagagacca      120
gggaaggggcc cccggtttgt gatgcgagtg ggcactggtg ggattgtggg atccaagggg      180
gatggcatcc ctgatcgctt ctcaagtctt ggctcaggcc tgaatcggtta cctgaccatc      240
aagaatatcc aggaagagga tgagagtgcac taccactgtg gggcagacca tggcagtggg      300
agcaacttcg tgtatgtctt cggaactggg accaagggtca ccgtccta      348

```

<210> SEQ ID NO 9
 <211> LENGTH: 116
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

Gln Pro Val Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala
1      5      10      15
Ser Val Thr Leu Thr Cys Thr Leu Ser Ser Gly Tyr Ser Asp Tyr Lys
      20      25      30
Val Asp Trp Tyr Gln Gln Arg Pro Gly Lys Gly Pro Arg Phe Val Met
      35      40      45
Arg Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Glu Gly Ile Pro
      50      55      60
Asp Arg Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Tyr Leu Thr Ile
      65      70      75      80
Lys Asn Ile Gln Glu Glu Asp Glu Ser Asp Tyr His Cys Gly Ala Asp
      85      90      95
His Gly Ser Gly Asn Asn Phe Val Tyr Val Phe Gly Thr Gly Thr Lys

```


-continued

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

```

Asp Ile Gln Leu Thr Pro Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ala Gly Trp
                20           25           30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
            35           40           45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
            50           55           60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asp Ser Phe Pro Pro
            85           90           95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
            100           105

```

<210> SEQ ID NO 14

<211> LENGTH: 321

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

```

gacatccagt tgaccccgtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtca gggatttgcc ggctgggttag cctgggtatca gcagaaacca    120
gggaaagccc ctaagctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca    180
agggttcagcg gcagtggatc tgggacagat ttcactctca ccatcagcag cctgcagcct    240
gaagattttg caacttacta ttgtcaacag gctgacagtt tccctccac tttcggcgga    300
gggaccaagg tggagatcaa a                                     321

```

<210> SEQ ID NO 15

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Val Ile Ser Ser Trp
                20           25           30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Ser Leu Leu Ile
            35           40           45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
            50           55           60
Ser Val Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65           70           75           80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe
            85           90           95
Thr Phe Gly Pro Gly Thr Lys Val Asp Phe Lys
            100           105

```

<210> SEQ ID NO 16

<211> LENGTH: 321

-continued

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

```

gacatccaga tgacccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtc ggttattagc agctggttag cctgggtatca gcagaaacca      120
gggaaagccc ctgacctctc gatctatgct gcatccagtt tgcaaagtgg ggtcccatca      180
aggttcagcg gcagtgtatc tgggacagat ttcactctca ccatcagcag cctgcagcct      240
gaagattttg caacttacta ttgtcaacag gctaacagtt tccattcac tttcgccct      300
gggaccaaag tggatttcaa a                                          321

```

<210> SEQ ID NO 17

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10          15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ser Ser Ser Trp
                20          25          30
Phe Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
                35          40          45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
                50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65          70          75          80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe
                85          90          95
Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
                100          105

```

<210> SEQ ID NO 18

<211> LENGTH: 321

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

```

gacatccaga tgacccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc      60
atcacttgtc gggcgagtc ggggaagtagc agctggtttg cctgggtatca gcagaaacca      120
gggaaagccc caaagctctc gatctatgct gcatccagtt tgcaaagtgg ggtcccatca      180
aggttcagcg gcagtggatc tgggacagac ttcactctca ccatcagcag cctgcagcct      240
gaagattttg caacttacta ttgtcaacag gctaacagtt tccattcac tttcgccct      300
gggaccaaag tggatatcaa a                                          321

```

<210> SEQ ID NO 19

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```

Asp Ser Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10          15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp
                20          25          30

```


-continued

Phe Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Asn Leu Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe
 85 90 95

Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
 100 105

<210> SEQ ID NO 20
 <211> LENGTH: 321
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

gacagccaga tgacccagtc tccatcttcc gtgtctgcct ctgtaggaga cagagtcacc	60
atcacttgtc gggcgagtc ggggtattagc agctgggttg cctgggtatca gcagaaacca	120
gggcaagccc ctaacctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca	180
aggttcagcg gcagtggatc tgggacagaa ttcactctca ccatcagcag cctgcagcct	240
gaagattttg caacttacta ttgtcaacag gctaacagtt tccattcac tttcggcct	300
gggaccaaag tggatatcaa a	321

<210> SEQ ID NO 21
 <211> LENGTH: 107
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Gly Gln Val Ile Ser Ser Trp
 20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Asp Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Thr Ser Phe Pro Leu
 85 90 95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105

<210> SEQ ID NO 22
 <211> LENGTH: 321
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

gacatccaga tgacccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc	60
atcacttgtc gggcggtgca ggttattagc agctgggttag cctgggtatca gcagaaacca	120
gggaaagccc ctaagctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatcg	180

-continued

```

aggttcagcg gcagtggatc tgggacagat ttcactctca ccatcagcag cctgcagcct 240
gacgattttg caacttacta ttgtcaacag gctaccagtt ttccctcac tttcggcgga 300
gggaccaagg tggagatcaa a 321

```

```

<210> SEQ ID NO 23
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 23

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Phe Ser Gly Trp
20          25          30
Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35          40          45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65          70          75          80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe
85          90          95
Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys
100         105

```

```

<210> SEQ ID NO 24
<211> LENGTH: 321
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 24

```

```

gacatccaga tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc 60
atcacttgtc gggcgagtca gggtttttagc ggttggttag cctggtatca gcagaaacca 120
gggaaagccc ctaagctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca 180
aggttcagtg gcagtggatc tgggacagat ttcactctca ccatcagcag cctgcagcct 240
gaagattttg caacttacta ctgtcaacag gctaacagtt tccattcac tttcggcct 300
gggaccaaag tggatatcaa a 321

```

```

<210> SEQ ID NO 25
<211> LENGTH: 107
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 25

```

```

Asp Ile Gln Leu Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1           5           10           15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Val Ile Ser Ser Trp
20          25          30
Phe Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Asn Leu Leu Ile
35          40          45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
50          55          60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65          70          75          80
Ala Asp Phe Ala Thr Tyr Phe Cys Gln Gln Ala Asn Ser Phe Pro Phe

```

-continued

85	90	95
Thr Phe Gly Pro Gly Thr Lys Val Asp Val Lys		
100	105	
 <210> SEQ ID NO 26		
<211> LENGTH: 321		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
 <400> SEQUENCE: 26		
gacatccagt tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc		60
atcacttgtc gggcgagtca gggtattagc agctggtttg cctgggtatca gcagaaacca		120
gggaaagccc ctaacctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca		180
aggttcagcg gcagtggatc tgggacagat ttcactctca ccatcagcag cctgcagcct		240
gcagattttg caacttactt ttgtcaacag gctaacagtt tccattcac ttctggccct		300
gggaccaaag tggatgtcaa a		321
 <210> SEQ ID NO 27		
<211> LENGTH: 107		
<212> TYPE: PRT		
<213> ORGANISM: Homo sapiens		
 <400> SEQUENCE: 27		
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly		
1	5	10 15
Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ser Ser Ser Trp		
20	25	30
Phe Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile		
35	40	45
Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly		
50	55	60
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
65	70	75 80
Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe		
85	90	95
Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys		
100	105	
 <210> SEQ ID NO 28		
<211> LENGTH: 321		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
 <400> SEQUENCE: 28		
gacatccaga tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc		60
atcacttgtc gggcgagtca gggtagtagc agctggtttg cctgggtatca acagaaacca		120
gggaaagccc caaagctcct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca		180
aggttcagcg gcagtggatc tgggacagat ttcactctca ccatcagcag cctgcagcct		240
gaagattttg caacttacta ttgtcaacag gctaacagtt tccattcac ttctggccct		300
gggaccaaag tggatatcaa a		321
 <210> SEQ ID NO 29		
<211> LENGTH: 107		
<212> TYPE: PRT		
<213> ORGANISM: Homo sapiens		

-continued

<400> SEQUENCE: 29

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Arg Asn Asp
 20 25 30

Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile
 35 40 45

Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser Tyr Pro Pro
 85 90 95

Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Glu
 100 105

<210> SEQ ID NO 30

<211> LENGTH: 108

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Consensus Sequence

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: Xaa can be Ile or Ser

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (4)..(4)

<223> OTHER INFORMATION: Xaa can be Met or Leu

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (29)..(29)

<223> OTHER INFORMATION: Xaa can be Gly or Val

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (30)..(30)

<223> OTHER INFORMATION: Xaa can be Ser, Phe or Ile

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (32)..(32)

<223> OTHER INFORMATION: Xaa can be Ser or Gly

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (34)..(34)

<223> OTHER INFORMATION: Xaa can be Phe or Leu

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (43)..(43)

<223> OTHER INFORMATION: Xaa can be Lys or Gln

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (46)..(46)

<223> OTHER INFORMATION: Xaa can be Lys, Asn or Ser

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (67)..(67)

<223> OTHER INFORMATION: Xaa can be Gly or Val

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (71)..(71)

<223> OTHER INFORMATION: Xaa can be Asp or Glu

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (82)..(82)

<223> OTHER INFORMATION: Xaa can be Glu or Ala

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (88)..(88)

<223> OTHER INFORMATION: Xaa can be Tyr or Phe

-continued

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (107)..(107)
<223> OTHER INFORMATION: Xaa can be Ile, Val or Phe

<400> SEQUENCE: 30

Asp Xaa Gln Xaa Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly
1             5             10             15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Xaa Xaa Ser Xaa
20             25             30

Trp Xaa Ala Trp Tyr Gln Gln Lys Pro Gly Xaa Ala Pro Xaa Leu Leu
35             40             45

Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser
50             55             60

Gly Ser Xaa Ser Gly Thr Xaa Phe Thr Leu Thr Ile Ser Ser Leu Gln
65             70             75             80

Pro Xaa Asp Phe Ala Thr Tyr Xaa Cys Gln Gln Ala Asn Ser Phe Pro
85             90             95

Phe Thr Phe Gly Pro Gly Thr Lys Val Asp Xaa Lys
100             105

<210> SEQ ID NO 31
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1             5             10             15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20             25             30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35             40             45

Ala Val Ile Trp Tyr Asp Gly Ser Asn Glu Tyr Tyr Ala Asp Ser Val
50             55             60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65             70             75             80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85             90             95

Ala Arg Asp Arg Gly Tyr Thr Ser Ser Trp Tyr Pro Asp Ala Phe Asp
100            105            110

Ile Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser
115            120

<210> SEQ ID NO 32
<211> LENGTH: 372
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

caggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc      60
tcctgtgcag cgtctggatt caccttcagt agctatggca tgcactgggt ccgccaggct      120
ccaggcaagg ggctggagtg ggtggcagtt atatggtatg atggaagtaa tgaatactat      180
gcagactccg tgaagggccg attcaccatc tccagagaca attccaagaa cacgctgtat      240
ctgcaaatga acagcctgag agccgaggac acggctgtgt attactgtgc gagagatcgg      300
gggtatacca gtagctggta ccctgatgct tttgatatct ggggccaaagg gacaatggtc      360

```

-continued

accgtctctt ca

372

<210> SEQ ID NO 33
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 33

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Arg Gly Tyr Ser Ser Ser Trp Tyr Pro Asp Ala Phe Asp
 100 105 110
 Ile Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 34
 <211> LENGTH: 121
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 34

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Val Ile Ser Phe Asp Gly Ser Leu Lys Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr Trp Gly
 100 105 110
 Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 35
 <211> LENGTH: 363
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 35

caggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
 tcctgtgcag cctctggatt caccttcagt agctatggca tgcactgggt ccgccaggct 120

-continued

```

ccaggcaagg ggctggagtg ggtggcagtt atatcatttg atggaagtct taaatactat 180
gcagactccg tgaagggccg attcaccatc tccagagaca attccaagaa caccctgtat 240
ctgcaaatga acagcctgag agctgaggac acggctgtgt attactgtgc gagagaacgg 300
actactttaa gtgggagcta ctttgactac tggggccagg gaaccctggt caccgtctcc 360
tca 363

```

```

<210> SEQ ID NO 36
<211> LENGTH: 121
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 36

```

```

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20          25          30
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Leu
35          40          45
Ser Val Ile Ser His Asp Gly Ser Ile Lys Tyr Tyr Ala Asp Ser Val
50          55          60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65          70          75          80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95
Ala Arg Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr Trp Gly
100         105         110
Gln Gly Thr Leu Val Thr Val Ser Ser
115         120

```

```

<210> SEQ ID NO 37
<211> LENGTH: 363
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 37

```

```

cagggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cctctggatt caccttcagt agctatgcca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg gttgtcagtt atatcacatg atggaagtat taaatactat 180
gcagactccg tgaagggccg attcaccatc tccagagaca attccaagaa caccgtgtat 240
ctgcaaatga acagcctgag agctgaggac acggctgtgt attactgtgc gagagaacgg 300
actactctaa gtgggagcta ctttgactac tggggccagg gaaccctggt caccgtctcc 360
tca 363

```

```

<210> SEQ ID NO 38
<211> LENGTH: 125
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 38

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20          25          30
Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

```

-continued

35	40	45	
Ser Tyr Ile Ser Ser Arg	Ser Ser Thr Ile Tyr	Ile Ala Asp Ser Val	
50	55	60	
Lys Gly Arg Phe Thr Ile	Ser Arg Asp Asn Ala	Lys Asn Ser Leu Tyr	
65	70	75	80
Leu Gln Met Asn Ser Leu	Arg Asp Glu Asp Thr	Ala Val Tyr Tyr Cys	
	85	90	95
Ala Arg Arg Ile Ala Ala	Ala Gly Gly Phe His	Tyr Tyr Tyr Ala Leu	
	100	105	110
Asp Val Trp Gly Gln Gly	Thr Thr Val Thr Val	Ser Ser	
	115	120	125

<210> SEQ ID NO 39
 <211> LENGTH: 375
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 39

gaggtgcagc tgggtggagtc tgggggaggc ctggtacagc ctgggggggtc cctgagactc	60
tcctgtgcag cctctggatt caccttcagt agctatagta tgaactgggt ccgccaggct	120
ccagggaagg ggctggagtg ggtttcgtac attagtagta ggagtagtac catatacatc	180
gcagactctg tgaaggggccg attcaccatc tccagagaca atgccaagaa ctcaactgtat	240
ctgcaaatga acagcctgag agacgaagac acggctgtgt attactgtgc gagacggata	300
gcagcagctg gtgggttcca ctactactac gctttggacg tctggggcca agggaccacg	360
gtcaccgtct cctca	375

<210> SEQ ID NO 40
 <211> LENGTH: 125
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 40

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1	15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr	
20	30
Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val	
35	45
Ser Tyr Ile Ser Ser Ser Ser Ser Thr Arg Tyr His Ala Asp Ser Val	
50	60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr	
65	80
Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys	
	95
Ala Arg Arg Ile Ala Ala Ala Gly Pro Trp Gly Tyr Tyr Tyr Ala Met	
100	110
Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
115	125

<210> SEQ ID NO 41
 <211> LENGTH: 375
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 41

-continued

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Thr Tyr
 20 25 30

Tyr Trp Ser Trp Ile Arg Gln Pro Ala Gly Lys Gly Leu Glu Trp Ile
 35 40 45

Gly Leu Ile Tyr Thr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Val Thr Met Ser Leu Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Arg Leu Thr Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala
 85 90 95

Arg Asp Arg Gly Tyr Tyr Tyr Gly Val Asp Val Trp Gly Gln Gly Thr
 100 105 110

Thr Val Thr Val Ser Ser
 115

<210> SEQ ID NO 45
 <211> LENGTH: 354
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

caggtgcagc tgcaggagtc gggcccagga ctggtgaagc ctcgggagac cctgtccctc	60
acctgcactg tctctggtgg ctccatcagt acttactact ggagctggat ccggcagccc	120
gccgggaagg gactggagtg gattgggctt atctatacca gtggggagcac caactacaac	180
ccctccctca agagtcgagt caccatgtca ttagacacgt ccaagaacca gttctccctg	240
aggctgacct ctgtgaccgc cgcggacacg gccgtttatt actgtgcgag agatcgtggg	300
tactactacg gtgtggacgt ctggggccag gggaccacgg tcaccgtctc ctca	354

<210> SEQ ID NO 46
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
 20 25 30

Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Ile Gly His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser
 50 55 60

Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe
 65 70 75 80

Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95

Cys Ala Lys Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 47
 <211> LENGTH: 360
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 47

```

caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc   60
acctgcactg tctctggtgg ctccatcagc agtgggtggtt actactggag ctggatccgc   120
cagcaccagc ggaagggcct ggagtggatt gggcacatcc attacagtgg gaacacctac   180
tacaaccctg ccctcaagag tcgagttacc atatcagtag acacgtctaa gaatcagttc   240
tccctgaaac tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgaaaaat   300
cgcggttctt actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca   360

```

<210> SEQ ID NO 48

<211> LENGTH: 120

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

```

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1           5           10          15
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Asn Ser Gly
          20          25          30
Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
          35          40          45
Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Ser Tyr Tyr Asn Pro Ser
          50          55          60
Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Gln Asn Gln Phe
65          70          75          80
Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
          85          90          95
Cys Ala Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln
          100         105         110
Gly Thr Thr Val Thr Val Ser Ser
          115          120

```

<210> SEQ ID NO 49

<211> LENGTH: 360

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

```

caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc   60
acctgcactg tctctggtgg ctccatcaac agtgggtggtt actactggag ctggatccgc   120
cagcaccagc ggaagggcct ggagtggatt gggtacatct attacagtgg gagctcctac   180
tacaaccctg ccctcaagag tcgagttacc atatcagtag acacgtctca gaaccagttc   240
tccctgaagc tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgagagat   300
cgggggcact actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca   360

```

<210> SEQ ID NO 50

<211> LENGTH: 120

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

```

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1           5           10          15
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
          20          25          30

```

-continued

Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser
 50 55 60

Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe
 65 70 75 80

Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95

Cys Ala Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 51
 <211> LENGTH: 360
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51

cagggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc	60
acctgcactg tctctggtgg ctccatcagt agtggtggtt actactggag ctggatccgc	120
cagcacccag ggaagggcct ggagtggatt gggtacattt attacagtgg gagcacctac	180
tacaaccctg ccctcaagag tcgagttacc atatcagtag acacgtctaa gaaccagttc	240
tccctgaagc tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgagagat	300
cggggccact actatggaat ggacgtctgg ggccaaggga ccacggtcac cgtctctca	360

<210> SEQ ID NO 52
 <211> LENGTH: 118
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 52

Gln Val Gln Leu Gln Glu Ser Gly Pro Arg Leu Val Lys Pro Ser Glu
 1 5 10 15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Asp Ser Ile Ser Ser Tyr
 20 25 30

Phe Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45

Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Val Thr Ile Ser Ile Asp Thr Ser Lys Asn Gln Phe Ser Leu
 65 70 75 80

Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Thr
 85 90 95

Arg Asp Arg Gly Ser Tyr Tyr Gly Ser Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Leu Val Thr Val Ser Ser
 115

<210> SEQ ID NO 53
 <211> LENGTH: 354
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

-continued

```

cagggtgcagc tgcaggagtc gggcccaaga ctggtgaagc cttcggagac cctgtccctc   60
acctgcactg tctctggtga ctccatcagt agttacttct ggagctggat ccggcagccc   120
ccagggaagg gactggagtg gcttgggtat atctattaca gtgggagcac caactacaac   180
ccctccctca agagtcgagt caccatatca atagacacgt ccaagaacca gttctccctg   240
aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt actgtacgag agatcggggg   300
agctactacg gatctgaact ctggggccag ggaacctggg tcacctctc ctca          354

```

```

<210> SEQ ID NO 54
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 54

```

```

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1           5           10           15
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
20          25          30
Gly Tyr Tyr Trp Thr Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
35          40          45
Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser
50          55          60
Leu Lys Ser Arg Ile Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe
65          70          75          80
Ser Leu Ser Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
85          90          95
Cys Ala Arg Asn Arg Gly Tyr Tyr Tyr Gly Met Asp Val Trp Gly Gln
100         105         110
Gly Thr Thr Val Thr Val Ser Ser
115          120

```

```

<210> SEQ ID NO 55
<211> LENGTH: 360
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 55

```

```

cagggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc   60
acctgcactg tctctggtgg ctccatcagc agtgggtggt actactggac ctggatccgc   120
cagcaccagc ggaagggcct ggagtggatt gggtagatct attacagtgg gaacacctac   180
tacaaccgct cctcaagag tcgaattacc atatcagtgg acacgtctaa gaaccagttc   240
tccctgagcc tgagctctgt gactgcccgc gacacggccg tgtattactg tgcgagaaat   300
cgcggttact actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca   360

```

```

<210> SEQ ID NO 56
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 56

```

```

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1           5           10           15
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly
20          25          30
Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu

```

-continued

35	40	45	
Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser			
50	55	60	
Leu Lys Ser Arg Val Thr Met Ser Val Asp Thr Ser Lys Asn Gln Phe			
65	70	75	80
Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr			
	85	90	95
Cys Ala Lys Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln			
	100	105	110
Gly Thr Thr Val Thr Val Ser Ser			
	115	120	
<210> SEQ ID NO 57			
<211> LENGTH: 360			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 57			
caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc			60
acctgcactg tctctggtgg ctccatcagc agtgggtggtt actactggag ctggatccgc			120
cagcaccagc ggaagggcct ggagtggatt gggtagatct attacagtgg gagcacctac			180
tacaaccctg cctcaagag tcgagttacc atgtcagtag acacgtctaa gaaccagttc			240
tccctgaaac tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgaaaaat			300
cgcggttctt actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctctctca			360
<210> SEQ ID NO 58			
<211> LENGTH: 120			
<212> TYPE: PRT			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 58			
Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln			
1	5	10	15
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Asn Ser Gly			
	20	25	30
Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu			
	35	40	45
Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Ser Tyr Tyr Asn Pro Ser			
50	55	60	
Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe			
65	70	75	80
Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr			
	85	90	95
Cys Ala Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln			
	100	105	110
Gly Thr Thr Val Thr Val Ser Ser			
	115	120	
<210> SEQ ID NO 59			
<211> LENGTH: 360			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 59			
caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc			60

-continued

```

acctgcactg tctctggtgg ctccatcaat agtgggtggtt actactggag ctggatccgc 120
cagcaccagc ggaagggcct ggagtggtt ggtacatct attacagtgg gagcagctac 180
tacaaccgct ccctcaagag tcgagttacc atatcagttg acacgtctaa gaaccagttc 240
tccctgaagc tgagttctgt gactgccgcg gacacggccg tgtattactg tgcgagagat 300
cgggggcact actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca 360

```

```

<210> SEQ ID NO 60
<211> LENGTH: 123
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 60

```

```

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1           5           10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
          20          25          30
Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
          35          40          45
Ala Leu Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val
          50          55          60
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
          65          70          75          80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
          85          90          95
Ala Arg Glu Asn Thr Val Thr Ile Tyr Tyr Asn Tyr Gly Met Asp Val
          100         105         110
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
          115         120

```

```

<210> SEQ ID NO 61
<211> LENGTH: 116
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa can be Val or Glu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: Xaa can be Asn or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (38)..(38)
<223> OTHER INFORMATION: Xaa can be Gln or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(55)
<223> OTHER INFORMATION: Xaa can be Ile or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (61)..(61)
<223> OTHER INFORMATION: Xaa can be Asp or Glu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (77)..(77)
<223> OTHER INFORMATION: Xaa can be Tyr or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (101)..(101)
<223> OTHER INFORMATION: Xaa can be Ser or Asn

```

```

<400> SEQUENCE: 61

```

-continued

Gln Pro Xaa Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala
 1 5 10 15
 Ser Val Thr Leu Thr Cys Thr Leu Xaa Ser Gly Tyr Ser Asp Tyr Lys
 20 25 30
 Val Asp Trp Tyr Gln Xaa Arg Pro Gly Lys Gly Pro Arg Phe Val Met
 35 40 45
 Arg Val Gly Thr Gly Gly Xaa Val Gly Ser Lys Gly Xaa Gly Ile Pro
 50 55 60
 Asp Arg Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Xaa Leu Thr Ile
 65 70 75 80
 Lys Asn Ile Gln Glu Glu Asp Glu Ser Asp Tyr His Cys Gly Ala Asp
 85 90 95
 His Gly Ser Gly Xaa Asn Phe Val Tyr Val Phe Gly Thr Gly Thr Lys
 100 105 110
 Val Thr Val Leu
 115

<210> SEQ ID NO 62
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

Thr Gly Ser Ser Ser Asn Thr Gly Ala Gly Tyr Asp Val His
 1 5 10

<210> SEQ ID NO 63
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

Gly Ser Gly Asn Arg Pro Ser
 1 5

<210> SEQ ID NO 64
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

Gln Ser Tyr Asp Ser Ser Leu Ser Gly Trp Val
 1 5 10

<210> SEQ ID NO 65
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Asp Val His
 1 5 10

<210> SEQ ID NO 66
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

Gly Ser Asn Asn Arg Pro Ser
 1 5

-continued

<210> SEQ ID NO 67
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

Met Ile Trp His Ser Ser Ala Ser Val
1 5

<210> SEQ ID NO 68
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

Thr Leu Arg Ser Gly Ile Asn Val Gly Thr Tyr Arg Ile Tyr
1 5 10

<210> SEQ ID NO 69
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

Tyr Lys Ser Asp Ser Asp Lys Gln Gln Gly Ser
1 5 10

<210> SEQ ID NO 70
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 70

Gly Ala Asp His Gly Ser Gly Ser Asn Phe Val Tyr Val
1 5 10

<210> SEQ ID NO 71
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 71

Thr Leu Asn Ser Gly Tyr Ser Asp Tyr Lys Val
1 5 10

<210> SEQ ID NO 72
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Asp
1 5 10

<210> SEQ ID NO 73
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

Gly Ala Asp His Gly Ser Gly Asn Asn Phe Val Tyr Val
1 5 10

-continued

<210> SEQ ID NO 74
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

Thr Leu Ser Ser Gly Tyr Ser Asp Tyr Lys Val
1 5 10

<210> SEQ ID NO 75
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Glu
1 5 10

<210> SEQ ID NO 76
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

Gln Gln Ala Asn Ser Phe Pro Phe Thr
1 5

<210> SEQ ID NO 77
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Arg Ala Ser Gln Gly Phe Ser Gly Trp Leu Ala
1 5 10

<210> SEQ ID NO 78
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 78

Val Gly Thr Gly Gly Thr Val Gly Ser Lys Gly Glu
1 5 10

<210> SEQ ID NO 79
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 79

Gln Gln Ala Thr Ser Phe Pro Leu Thr
1 5

<210> SEQ ID NO 80
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 80

Arg Ala Ser Gln Val Ile Ser Ser Trp Leu Ala
1 5 10

<210> SEQ ID NO 81
<211> LENGTH: 7

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 81

Ala Ala Ser Ser Leu Gln Ser
1 5

<210> SEQ ID NO 82
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 82

Gln Gln Ala Asp Ser Phe Pro Pro Thr
1 5

<210> SEQ ID NO 83
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

Arg Ala Ser Gln Val Ile Ser Ser Trp Phe Ala
1 5 10

<210> SEQ ID NO 84
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

Leu Gln His Asn Ser Tyr Pro Pro Thr
1 5

<210> SEQ ID NO 85
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 85

Arg Ala Ser Gln Gly Ser Ser Ser Trp Phe Ala
1 5 10

<210> SEQ ID NO 86
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 86

Arg Ala Ser Gln Gly Ile Ser Ser Trp Phe Ala
1 5 10

<210> SEQ ID NO 87
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 87

Arg Ala Gly Gln Val Ile Ser Ser Trp Leu Ala
1 5 10

<210> SEQ ID NO 88
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 88

Arg Ala Ser Gln Gly Ile Ala Gly Trp Leu Ala
1 5 10

<210> SEQ ID NO 89

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Arg Ala Ser Gln Gly Ile Arg Asn Asp Leu Gly
1 5 10

<210> SEQ ID NO 90

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Leu Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 91

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Ser Tyr Gly Met His
1 5

<210> SEQ ID NO 92

<211> LENGTH: 17

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Val Ile Trp Tyr Asp Gly Ser Asn Glu Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 93

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Asp Arg Gly Tyr Thr Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile
1 5 10 15

<210> SEQ ID NO 94

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94

Ser Tyr Ala Met His
1 5

<210> SEQ ID NO 95

<211> LENGTH: 17

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 95

Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys
1           5              10              15

Gly

<210> SEQ ID NO 96
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 96

Asp Arg Gly Tyr Ser Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile
1           5              10              15

<210> SEQ ID NO 97
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97

Thr Tyr Ser Met Asn
1           5

<210> SEQ ID NO 98
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

Val Ile Ser Phe Asp Gly Ser Leu Lys Tyr Tyr Ala Asp Ser Val Lys
1           5              10              15

Gly

<210> SEQ ID NO 99
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr
1           5              10

<210> SEQ ID NO 100
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 100

Ser Tyr Ser Met Asn
1           5

<210> SEQ ID NO 101
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 101

Val Ile Ser His Asp Gly Ser Ile Lys Tyr Tyr Ala Asp Ser Val Lys
1           5              10              15

Gly

```

-continued

<210> SEQ ID NO 102
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

Arg Ile Ala Ala Ala Gly Gly Phe His Tyr Tyr Tyr Ala Leu Asp Val
1 5 10 15

<210> SEQ ID NO 103
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

Ser Phe Ser Met Asn
1 5

<210> SEQ ID NO 104
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 104

Tyr Ile Ser Ser Arg Ser Ser Thr Ile Tyr Ile Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 105
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

Arg Ile Ala Ala Ala Gly Pro Trp Gly Tyr Tyr Tyr Ala Met Asp Val
1 5 10 15

<210> SEQ ID NO 106
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 106

Ser Gly Gly Tyr Tyr Trp Thr
1 5

<210> SEQ ID NO 107
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 107

Tyr Ile Ser Ser Ser Ser Ser Thr Arg Tyr His Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 108
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108

<400> SEQUENCE: 115

-continued

Thr Tyr Tyr Trp Ser
1 5

<210> SEQ ID NO 116
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 116

His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 117
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 117

Asp Arg Gly Ser Tyr Tyr Gly Ser Asp Tyr
1 5 10

<210> SEQ ID NO 118
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 118

Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 119
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 119

Asp Arg Gly Tyr Tyr Tyr Gly Val Asp Val
1 5 10

<210> SEQ ID NO 120
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 120

Tyr Ile Tyr Tyr Ser Gly Ser Ser Tyr Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 121
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 121

Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 122
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <400> SEQUENCE: 122

Leu Ile Tyr Thr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

-continued

<210> SEQ ID NO 123
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa can be Gly or Val
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa can be Ile, Phe or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa can be Ser or Gly
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (10)..(10)
<223> OTHER INFORMATION: Xaa can be Phe or Leu.

<400> SEQUENCE: 123

Arg Ala Ser Gln Xaa Xaa Ser Xaa Trp Xaa Ala
1 5 10

<210> SEQ ID NO 124
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa can be Asn or Ser

<400> SEQUENCE: 124

Thr Leu Xaa Ser Gly Tyr Ser Asp Tyr Lys Val Asp
1 5 10

<210> SEQ ID NO 125
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa can be Ile or Thr

<400> SEQUENCE: 125

Thr Gly Ser Ser Ser Asn Xaa Gly Ala Gly Tyr Asp Val His
1 5 10

<210> SEQ ID NO 126
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa can be Ile or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: Xaa can be Asp or Glu

-continued

<400> SEQUENCE: 126

Val	Gly	Thr	Gly	Gly	Xaa	Val	Gly	Ser	Lys	Gly	Xaa
1			5						10		

<210> SEQ ID NO 127

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Consensus sequence

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: Xaa can be Asn or Gly

<400> SEQUENCE: 127

Gly	Ser	Xaa	Asn	Arg	Pro	Ser
1			5			

<210> SEQ ID NO 128

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Consensus sequence

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (8)..(8)

<223> OTHER INFORMATION: Xaa can be Ser or Asn

<400> SEQUENCE: 128

Gly	Ala	Asp	His	Gly	Ser	Gly	Xaa	Asn	Phe	Val	Tyr	Val
1				5					10			

<210> SEQ ID NO 129

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Consensus sequence

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: Xaa can be Ser or Thr

<400> SEQUENCE: 129

Ser	Gly	Gly	Tyr	Tyr	Trp	Xaa
1			5			

<210> SEQ ID NO 130

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Consensus sequence

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: Xaa can be Gly or Ala

<400> SEQUENCE: 130

Ser	Tyr	Xaa	Met	His
1			5	

<210> SEQ ID NO 131

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Artificial

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa can be Ser or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa can be Ser or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa can be Tyr or Phe

<400> SEQUENCE: 131

Xaa Xaa Ser Met Asn
1 5

<210> SEQ ID NO 132
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa can be Tyr or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa can be Thr or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa can be Ser or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa can be Thr or Ser

<400> SEQUENCE: 132

Xaa Ile Xaa Tyr Ser Gly Xaa Xaa Tyr Tyr Asn Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 133
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa can be Phe or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa can be Leu or Thr

<400> SEQUENCE: 133

Val Ile Ser Xaa Asp Gly Ser Xaa Lys Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> SEQ ID NO 134
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence

-continued

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa can be Arg or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa can be Ile or Arg
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (11)..(11)
<223> OTHER INFORMATION: Xaa can be Ile, His or Try

```

```

<400> SEQUENCE: 134

```

```

Tyr Ile Ser Ser Xaa Ser Ser Thr Xaa Tyr Xaa Ala Asp Ser Val Lys
1           5              10              15

```

```

Gly

```

```

<210> SEQ ID NO 135
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa can be Lys or Glu

```

```

<400> SEQUENCE: 135

```

```

Val Ile Trp Tyr Asp Gly Ser Asn Xaa Tyr Tyr Ala Asp Ser Val Lys
1           5              10              15

```

```

Gly

```

```

<210> SEQ ID NO 136
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa can be Asn or Asp
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa can be His, Tyr or Phe

```

```

<400> SEQUENCE: 136

```

```

Xaa Arg Gly Xaa Tyr Tyr Gly Met Asp Val
1           5              10

```

```

<210> SEQ ID NO 137
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa can be Gly or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Xaa can be Phe or Trp
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: Xaa can be His or Gly

```

-continued

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Xaa can be Leu and Met

<400> SEQUENCE: 137

Arg Ile Ala Ala Ala Gly Xaa Xaa Xaa Tyr Tyr Tyr Ala Xaa Asp Val
1             5             10             15

```

```

<210> SEQ ID NO 138
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa can be Ser or Thr

<400> SEQUENCE: 138

```

```

Asp Arg Gly Tyr Xaa Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile
1             5             10             15

```

```

<210> SEQ ID NO 139
<211> LENGTH: 112
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (29)..(29)
<223> OTHER INFORMATION: Xaa can be Ile or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (41)..(41)
<223> OTHER INFORMATION: Xaa can be Val or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(54)
<223> OTHER INFORMATION: Xaa can be Gly or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (107)..(107)
<223> OTHER INFORMATION: Xaa can be Arg or Lys

```

```

<400> SEQUENCE: 139

Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
1             5             10             15

Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Xaa Gly Ala Gly
20            25            30

Tyr Asp Val His Trp Tyr Gln Gln Xaa Pro Gly Thr Ala Pro Lys Leu
35            40            45

Leu Ile Tyr Gly Ser Xaa Asn Arg Pro Ser Gly Val Pro Asp Arg Phe
50            55            60

Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu
65            70            75            80

Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser
85            90            95

Leu Ser Gly Trp Val Phe Gly Gly Gly Thr Xaa Arg Leu Thr Val Leu
100           105           110

```

```

<210> SEQ ID NO 140
<211> LENGTH: 120
<212> TYPE: PRT
<213> ORGANISM: Artificial

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: Xaa can be And or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (37)..(37)
<223> OTHER INFORMATION: Xaa can be Ser or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (52)..(52)
<223> OTHER INFORMATION: Xaa can be Tyr or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(54)
<223> OTHER INFORMATION: Xaa can be Tyr or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa can be Ser or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (59)..(59)
<223> OTHER INFORMATION: Xaa can be Ser or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (69)..(69)
<223> OTHER INFORMATION: Xaa can be Ser or Thr
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (71)..(71)
<223> OTHER INFORMATION: Xaa can be Val or Ile
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (77)..(77)
<223> OTHER INFORMATION: Xaa can be Ile or Met
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (83)..(83)
<223> OTHER INFORMATION: Xaa can be Lys or Gln
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(99)
<223> OTHER INFORMATION: Xaa can be Arg or Lys
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (100)..(100)
<223> OTHER INFORMATION: Xaa can be Asp or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (103)..(103)
<223> OTHER INFORMATION: Xaa can be His, Phe or Try

<400> SEQUENCE: 140

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1          5          10          15

Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Xaa Ser Gly
20          25          30

Gly Tyr Tyr Trp Xaa Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
35          40          45

Trp Ile Gly Xaa Ile Xaa Tyr Ser Gly Xaa Xaa Tyr Tyr Asn Pro Ser
50          55          60

Leu Lys Ser Arg Xaa Thr Xaa Ser Val Asp Thr Ser Xaa Asn Gln Phe
65          70          75          80

Ser Leu Xaa Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
85          90          95

Cys Ala Xaa Xaa Arg Gly Xaa Tyr Tyr Gly Met Asp Val Trp Gly Gln
100         105         110

Gly Thr Thr Val Thr Val Ser Ser

```

-continued

115

120

```

<210> SEQ ID NO 141
<211> LENGTH: 125
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (23)..(23)
<223> OTHER INFORMATION: Xaa can be Ala or Val
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(24)
<223> OTHER INFORMATION: Xaa can be Ala or Val
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(31)
<223> OTHER INFORMATION: Xaa can be Thr or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (32)..(32)
<223> OTHER INFORMATION: Xaa can be Tyr or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(54)
<223> OTHER INFORMATION: Xaa can be Ser or Arg
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa can be Arg or Ile
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (60)..(60)
<223> OTHER INFORMATION: Xaa can be His, Try or Ile
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (105)..(105)
<223> OTHER INFORMATION: Xaa can be Pro or Gly
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (106)..(106)
<223> OTHER INFORMATION: Xaa can be Trp or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (107)..(107)
<223> OTHER INFORMATION: Xaa can be Gly or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (112)..(112)
<223> OTHER INFORMATION: Xaa can be Met or Leu

<400> SEQUENCE: 141

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1          5          10          15

Ser Leu Arg Leu Ser Cys Xaa Xaa Ser Gly Phe Thr Phe Ser Xaa Xaa
20        25        30

Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35        40        45

Ser Tyr Ile Ser Ser Xaa Ser Ser Thr Xaa Tyr Xaa Ala Asp Ser Val
50        55        60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65        70        75        80

Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys
85        90        95

Ala Arg Arg Ile Ala Ala Ala Gly Xaa Xaa Xaa Tyr Tyr Tyr Ala Xaa
100       105       110

Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
115       120       125

```

-continued

```

<210> SEQ ID NO 142
<211> LENGTH: 121
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (33)..(33)
<223> OTHER INFORMATION: Xaa can be Gly or Ala
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (48)..(48)
<223> OTHER INFORMATION: Xaa can be Val or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (49)..(49)
<223> OTHER INFORMATION: Xaa can be Ala or Ser
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (53)..(53)
<223> OTHER INFORMATION: Xaa can be Phe or His
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (57)..(57)
<223> OTHER INFORMATION: Xaa can be Leu or Ile

<400> SEQUENCE: 142

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1             5             10             15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20             25             30

Xaa Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Xaa
35             40             45

Xaa Val Ile Ser Xaa Asp Gly Ser Xaa Lys Tyr Tyr Ala Asp Ser Val
50             55             60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
65             70             75             80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85             90             95

Ala Arg Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr Trp Gly
100            105            110

Gln Gly Thr Leu Val Thr Val Ser Ser
115            120

```

```

<210> SEQ ID NO 143
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Consensus Sequence
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (58)..(58)
<223> OTHER INFORMATION: Xaa can be Glu or Lys
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (103)..(103)
<223> OTHER INFORMATION: Xaa can be Thr or Ser

<400> SEQUENCE: 143

Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg
1             5             10             15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20             25             30

```


-continued

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ala Val Ile Trp Tyr Asp Gly Ser Asn Xaa Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Asp Arg Gly Tyr Xaa Ser Ser Trp Tyr Pro Asp Ala Phe Asp
 100 105 110

Ile Trp Gly Gln Gly Thr Met Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 144
 <211> LENGTH: 1026
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 144

```

aactcgggtga acaactgagg gaaccaaacc agagacgcgc tgaacagaga gaatcaggct      60
caaagcaagt ggaagtgggc agagattcca ccaggactgg tgcaaggcgc agagccagcc      120
agatttgaga agaaggcaaa aagatgctgg ggagcagagc tgtaatgctg ctgttgctgc      180
tgccctggac agctcagggc agagctgtgc ctgggggcag cagccctgcc tggactcagt      240
gccagcagct ttcacagaag ctctgcacac tggcctggag tgcacatcca ctagtgggac      300
acatggatct aagagaagag ggagatgaag agactacaaa tgatgttccc catatccagt      360
gtggagatgg ctgtgacccc caaggactca gggacaacag tcagttctgc ttgcaaagga      420
tccaccaggg tctgattttt tatgagaagc tgctaggatc ggatattttc acagggggagc      480
cttctctgct ccctgatagc cctgtgggcc agcttcatgc ctccctactg ggccctcagcc      540
aactcctgca gcctgagggt caccactggg agactcagca gattocaagc ctcagtccca      600
gccagccatg gcagcgtctc cttctccgct tcaaaatcct tcgcagcctc caggcctttg      660
tggtctgtagc cggccgggtc ttgcccattg gagcagcaac cctgagtccc taaaggcagc      720
agctcaagga tggcactcag atctccatgg ccagcaagg ccaagataaa tctaccaccc      780
caggcacctg tgagccaaca ggttaattag tccattaatt ttagtgggac ctgcatatgt      840
tgaaaattac caatactgac tgacatgtga tgctgaccta tgataagggt gagtatttat      900
tagatgggaa gggaaatttg gggattatct atcctcctgg ggacagtttg gggaggatta      960
tttattgtat ttatattgaa ttatgtactt ttttcaataa agtcttattt ttgtggctaa     1020
aaaaaa                                           1026

```

<210> SEQ ID NO 145
 <211> LENGTH: 189
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 145

Met Leu Gly Ser Arg Ala Val Met Leu Leu Leu Leu Leu Pro Trp Thr
 1 5 10 15

Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser Pro Ala Trp Thr Gln
 20 25 30

Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His
 35 40 45

-continued

Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr
50 55 60

Thr Asn Asp Val Pro His Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln
65 70 75 80

Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln Arg Ile His Gln Gly
85 90 95

Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu
100 105 110

Pro Ser Leu Leu Pro Asp Ser Pro Val Gly Gln Leu His Ala Ser Leu
115 120 125

Leu Gly Leu Ser Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr
130 135 140

Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln Pro Trp Gln Arg Leu Leu
145 150 155 160

Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val Ala
165 170 175

Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro
180 185

<210> SEQ ID NO 146

<211> LENGTH: 1399

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 146

```

ctgtttcagg gccattggac tctccgtcct gccagagca agatgtgtca ccagcagttg      60
gtcatctctt ggtttttcct ggtttttctg gcatctcccc tcgtggccat atgggaactg      120
aagaaagatg tttatgtcgt agaattggat tggatccgg atgcccctgg agaaatggtg      180
gtcctcacct gtgacacccc tgaagaagat ggtatcacct ggaccttga ccagagcagt      240
gaggtcttag gctctggcaa aacctgacc atccaagtca aagagtttg agatgctggc      300
cagtacacct gtcacaaagg aggcgagggt ctaagccatt cgctcctgct gcttcacaaa      360
aaggaagatg gaatttggtc cactgatatt ttaaaggacc agaaagaacc caaaaataag      420
acctttctaa gatcgaggc caagaattat tctggacgtt tcacctgctg gtggctgacg      480
acaatcagta ctgatttgac attcagtgtc aaaagcagca gaggtcttct tgaccccaaa      540
ggggtgacgt gcgagctgc tacactctct gcagagagag tcagagggga caacaaggag      600
tatgagtact cagtggagtg ccaggaggac agtgccctgcc cagctgctga ggagagtctg      660
cccattgagg tcattggtga tgcggtcac aagctcaagt atgaaaacta caccagcagc      720
ttcttcatca gggacatcat caaacctgac ccaccaaga acttgacgct gaagccatta      780
aagaattctc ggcaggtgga ggtcagctgg gagtaccctg acacctggag tactccacat      840
tcctacttct ccctgacatt ctgcgttcag gtccagggca agagcaagag agaaaagaaa      900
gatagagtct tcacggacaa gacctcagcc acggctcatct gccgcaaaaa tgccagcatt      960
agcgtgcggg cccaggaccg ctactatagc tcactttgga gcgaatgggc atctgtgccc     1020
tgcagttagg ttctgatcca ggatgaaaat ttggaggaaa agtggaagat attagcaaaa     1080
atgtttaaag acacaacgga atagacccaa aaagataatt tctatctgat ttgctttaaa     1140
acgttttttt aggatcacia tgatatcttt gctgtatttg tatagttaga tgctaaatgc     1200
tcattgaaac aatcagctaa tttatgtata gattttccag ctctcaagtt gccatggggc     1260
ttcatgctat ttaaatatth aagtaattta tgtatttatt agtatattac tgttatttaa     1320

```

-continued

```
cgtttgctcg ccaggatgta tggaatgttt cataactctta tgacctgatc catcaggatc 1380
agtcacctatt atgcaaaat 1399
```

```
<210> SEQ ID NO 147
<211> LENGTH: 328
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 147
```

```
Met Cys His Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Phe Leu
 1          5          10          15

Ala Ser Pro Leu Val Ala Ile Trp Glu Leu Lys Lys Asp Val Tyr Val
 20          25          30

Val Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu
 35          40          45

Thr Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln
 50          55          60

Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
 65          70          75          80

Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
 85          90          95

Leu Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp
100          105          110

Ser Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe
115          120          125

Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
130          135          140

Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg
145          150          155          160

Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser
165          170          175

Ala Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu
180          185          190

Cys Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile
195          200          205

Glu Val Met Val Asp Ala Val His Lys Leu Lys Tyr Glu Asn Tyr Thr
210          215          220

Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn
225          230          235          240

Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp
245          250          255

Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr
260          265          270

Phe Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg
275          280          285

Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys Asn Ala
290          295          300

Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser
305          310          315          320

Glu Trp Ala Ser Val Pro Cys Ser
325
```

```
<210> SEQ ID NO 148
<211> LENGTH: 2826
<212> TYPE: DNA
```

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 148

```

acaaggggtg cagcctggct ctgaagtgga attatgtgct tcaaacaggt tgaagaggg      60
aaacagtctt ttctgtcttc cagacatgaa tcaggtcact attcaatggg atgcagtaat    120
agccctttac atactcttca gctggtgtca tggaggaatt acaaatataa actgctctgg    180
ccacatctgg gtagaaccag ccacaatttt taagatgggt atgaatatct ctatatattg    240
ccaagcagca attaagaact gccaaccaag gaaacttcat tttataaaa atggcatcaa    300
agaaagattt caaatcacia ggattaataa aacaacagct cggctttggg ataaaaactt    360
tctggaacca catgcttcta tgtactgcac tgctgaatgt cccaaacatt ttcaagagac    420
actgatatgt ggaaaagaca tttcttctgg atatccgcca gatattcctg atgaagtaac    480
ctgtgtcatt tatgaatatt caggcaacat gacttgcacc tggaatgctg ggaagctcac    540
ctacatagac acaaaatcac tggtacatgt gaagagttaa gagacagaag aagagcaaca    600
gtatctcacc tcaagctata ttaacatctc cactgattca ttacaagggt gcaagaagta    660
cttggtttgg gtccaagcag caaacgcact aggcattgaa gagtcaaac aactgcaaat    720
tcacctggat gatatagtga taccttctgc agccgtcatt tccagggctg agactataaa    780
tgctacagtg cccaagacca taatttattg ggatagtcaa acaacaattg aaaaggtttc    840
ctgtgaaatg agatacaagg ctacaacaaa ccaaacttgg aatgttaaag aatttgacac    900
caattttaca tatgtgcaac agtcagaatt ctacttgag ccaaacatta agtacgtatt    960
tcaagtgaga tgtcaagaaa caggcaaaag gtactggcag ccttgaggtt cactgttttt   1020
tcataaaaaca cctgaaacag ttccccaggt cacatcaaaa gcattccaac atgacacatg   1080
gaattctggg ctaacagttg ctccatctc tacagggcac cttacttctg acaacagagg   1140
agacattgga cttttatttg gaatgatcgt ctttgcgttt atgttgtcaa ttctttcttt   1200
gattgggata tttaacagat cattccgaac tgggattaaa agaaggatct tattgttaat   1260
accaaagtgg ctttatgaag atattcctaa tatgaaaaac agcaatgttg tgaaaatgct   1320
acaggaaaat agtgaactta tgaataataa ttccagttag caggtcctat atgttgatcc   1380
catgattaca gagataaaag aaatcttcat ccagaacac aagcctacag actacaagaa   1440
ggagaatata ggacccttgg agacaagaga ctaccgcga aactcgctat tcgacaatac   1500
tacagttgta tatattcctg atctcaacac tggatataaa ccccaaattt caaattttct   1560
gcctgagggg agccatctca gcaataataa tgaaattact tccttaacac ttaaacacc   1620
agttgattcc ttagactcag gaaataatcc caggttacaa aagcatccta attttgcttt   1680
ttctgtttca agtggtgaatt cactaagcaa cacaatattt cttggagaat taagcctcat   1740
attaaatcaa ggagaatgca gttctcctga catacaaac tcagtagagg aggaaaccac   1800
catgcttttg gaaaatgatt caccagtgga aactattcca gaacagacc tgcttctgga   1860
tgaatttgtc tcctgttttg ggatcgtgaa tgaggagttg ccatctatta atacttattt   1920
tccacaaaat attttggaag gccactcaa taggatttca ctcttggaag agtagagctg   1980
tgtggtcaaa atcaatatga gaaagctgcc ttgcaatctg aacttgggtt ttccctgcaa   2040
tagaaattga attctgcctc tttttgaaaa aaatgtattc acatacaaat ctccacatgg   2100
acacatgttt tcatttcctt tggataaata cctaggtagg ggattgctgg gccatatgat   2160
aagcatatgt ttcagttcta ccaatcttgt ttccagagta gtgacatttc tgtgtccta   2220
ccatcaccat gtaagaattc ccgggagctc catgcctttt taatttttag cattcttctg   2280

```

-continued

```

cctcatttct taaaattaga gaattaaggt cccgaagggt gaacatgctt catggtcaca 2340
catacaggca caaaaacagc attatgtgga cgcctcatgt attttttata gagtcaacta 2400
tttctctttt attttccctc attgaaagat gcaaaacagc tctctattgt gtacagaaag 2460
ggtaataaat gcaaaatacc tggtagtaaa ataaatgctg aaaattttcc tttaaaaatg 2520
aatcattagg ccaggcgtgg tggctcatgc ttgtaatccc agcactttgg taggctgagg 2580
taggtggatc acctgaggtc aggagttcga gtccagcctg gccaatatgc tgaaccctg 2640
tctctactaa aattacaaaa attagccggc catggtggca ggtgcttgta atcccagcta 2700
cttgggaggg tgaggcagga gaatcacttg aaccaggaag gcagaggttg cactgagctg 2760
agattgtgcc actgcactcc agcctgggca acaagagcaa aactctgtct ggaaaaaaaa 2820
aaaaaa 2826

```

<210> SEQ ID NO 149

<211> LENGTH: 629

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 149

```

Met Asn Gln Val Thr Ile Gln Trp Asp Ala Val Ile Ala Leu Tyr Ile
1           5           10          15

Leu Phe Ser Trp Cys His Gly Gly Ile Thr Asn Ile Asn Cys Ser Gly
20          25          30

His Ile Trp Val Glu Pro Ala Thr Ile Phe Lys Met Gly Met Asn Ile
35          40          45

Ser Ile Tyr Cys Gln Ala Ala Ile Lys Asn Cys Gln Pro Arg Lys Leu
50          55          60

His Phe Tyr Lys Asn Gly Ile Lys Glu Arg Phe Gln Ile Thr Arg Ile
65          70          75          80

Asn Lys Thr Thr Ala Arg Leu Trp Tyr Lys Asn Phe Leu Glu Pro His
85          90          95

Ala Ser Met Tyr Cys Thr Ala Glu Cys Pro Lys His Phe Gln Glu Thr
100         105         110

Leu Ile Cys Gly Lys Asp Ile Ser Ser Gly Tyr Pro Pro Asp Ile Pro
115         120         125

Asp Glu Val Thr Cys Val Ile Tyr Glu Tyr Ser Gly Asn Met Thr Cys
130         135         140

Thr Trp Asn Ala Gly Lys Leu Thr Tyr Ile Asp Thr Lys Tyr Val Val
145         150         155         160

His Val Lys Ser Leu Glu Thr Glu Glu Gln Gln Tyr Leu Thr Ser
165         170         175

Ser Tyr Ile Asn Ile Ser Thr Asp Ser Leu Gln Gly Gly Lys Lys Tyr
180         185         190

Leu Val Trp Val Gln Ala Ala Asn Ala Leu Gly Met Glu Glu Ser Lys
195         200         205

Gln Leu Gln Ile His Leu Asp Asp Ile Val Ile Pro Ser Ala Ala Val
210         215         220

Ile Ser Arg Ala Glu Thr Ile Asn Ala Thr Val Pro Lys Thr Ile Ile
225         230         235         240

Tyr Trp Asp Ser Gln Thr Thr Ile Glu Lys Val Ser Cys Glu Met Arg
245         250         255

Tyr Lys Ala Thr Thr Asn Gln Thr Trp Asn Val Lys Glu Phe Asp Thr
260         265         270

```

-continued

Asn Phe Thr Tyr Val Gln Gln Ser Glu Phe Tyr Leu Glu Pro Asn Ile
 275 280 285
 Lys Tyr Val Phe Gln Val Arg Cys Gln Glu Thr Gly Lys Arg Tyr Trp
 290 295 300
 Gln Pro Trp Ser Ser Leu Phe Phe His Lys Thr Pro Glu Thr Val Pro
 305 310 315 320
 Gln Val Thr Ser Lys Ala Phe Gln His Asp Thr Trp Asn Ser Gly Leu
 325 330 335
 Thr Val Ala Ser Ile Ser Thr Gly His Leu Thr Ser Asp Asn Arg Gly
 340 345 350
 Asp Ile Gly Leu Leu Leu Gly Met Ile Val Phe Ala Val Met Leu Ser
 355 360 365
 Ile Leu Ser Leu Ile Gly Ile Phe Asn Arg Ser Phe Arg Thr Gly Ile
 370 375 380
 Lys Arg Arg Ile Leu Leu Leu Ile Pro Lys Trp Leu Tyr Glu Asp Ile
 385 390 395 400
 Pro Asn Met Lys Asn Ser Asn Val Val Lys Met Leu Gln Glu Asn Ser
 405 410 415
 Glu Leu Met Asn Asn Asn Ser Ser Glu Gln Val Leu Tyr Val Asp Pro
 420 425 430
 Met Ile Thr Glu Ile Lys Glu Ile Phe Ile Pro Glu His Lys Pro Thr
 435 440 445
 Asp Tyr Lys Lys Glu Asn Thr Gly Pro Leu Glu Thr Arg Asp Tyr Pro
 450 455 460
 Gln Asn Ser Leu Phe Asp Asn Thr Thr Val Val Tyr Ile Pro Asp Leu
 465 470 475 480
 Asn Thr Gly Tyr Lys Pro Gln Ile Ser Asn Phe Leu Pro Glu Gly Ser
 485 490 495
 His Leu Ser Asn Asn Asn Glu Ile Thr Ser Leu Thr Leu Lys Pro Pro
 500 505 510
 Val Asp Ser Leu Asp Ser Gly Asn Asn Pro Arg Leu Gln Lys His Pro
 515 520 525
 Asn Phe Ala Phe Ser Val Ser Ser Val Asn Ser Leu Ser Asn Thr Ile
 530 535 540
 Phe Leu Gly Glu Leu Ser Leu Ile Leu Asn Gln Gly Glu Cys Ser Ser
 545 550 555 560
 Pro Asp Ile Gln Asn Ser Val Glu Glu Glu Thr Thr Met Leu Leu Glu
 565 570 575
 Asn Asp Ser Pro Ser Glu Thr Ile Pro Glu Gln Thr Leu Leu Pro Asp
 580 585 590
 Glu Phe Val Ser Cys Leu Gly Ile Val Asn Glu Glu Leu Pro Ser Ile
 595 600 605
 Asn Thr Tyr Phe Pro Gln Asn Ile Leu Glu Ser His Phe Asn Arg Ile
 610 615 620
 Ser Leu Leu Glu Lys
 625

<210> SEQ ID NO 150

<211> LENGTH: 2100

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 150

ggtaggctgaa cctgcagagt ggcagagagg ctccctctggg gctgtggggc tctacgtgga

60

-continued

tccgatggag	ccgctggtga	cctgggtggt	ccccctctc	tctctcttc	tgtgtccag	120
gcagggcgct	gcctgcagaa	ccagtgagt	ctgttttcag	gacccgccat	atccggatgc	180
agactcaggc	tcggcctcgg	gccctaggga	cctgagatgc	tatcggatat	ccagtgatcg	240
ttacgagtgc	tcctggcagt	atgagggtcc	cacagctggg	gtcagccact	tctgcggtg	300
ttgccttagc	tccgggcgct	gctgctactt	cgccgccggc	tcagccacca	ggctgcagtt	360
ctccgaccag	gctggggtgt	ctgtgctgta	cactgtcaca	ctctgggtgg	aatcctgggc	420
caggaaccag	acagagaagt	ctcctgaggt	gacctgcag	ctctacaact	cagttaaata	480
tgagcctcct	ctgggagaca	tcaaggtgtc	caagttggcc	gggcagctgc	gtatggagt	540
ggagaccccg	gataaccagg	ttggtgctga	ggtgcagttc	cggcacccga	caccagcag	600
cccatggaag	ttgggcgact	gcggaacctc	ggatgatgat	actgagtctc	gcctctgccc	660
cctggagatg	aatgtggccc	aggaattcca	gctccgacga	cggcagctgg	ggagccaagg	720
aagttcctgg	agcaagtgga	gcagccccgt	gtgcgttccc	cctgaaaacc	ccccacagcc	780
tcagtgagag	ttctcggtgg	agcagctggg	ccaggatggg	aggaggcggc	tgacctgaa	840
agagcagcca	accagctgg	agcttcagga	aggctgtcaa	gggctggcgc	ctggcacgga	900
ggtcacttac	cgactacagc	tccacatgct	gtcctgcccc	tgtaaggcca	aggccaccag	960
gacctgcac	ctggggaaga	tgcctatct	ctcggtgct	gcctacaacg	tggtgtcat	1020
ctcctcgaac	caatttggtc	ctggcctgaa	ccagacgtgg	cacattcctg	ccgacacca	1080
cacagaacca	gtggtcttga	atatcagcgt	cggaaaccaac	gggaccacca	tgtattggcc	1140
agcccggtct	cagagcatga	cgtattgcat	tgaatggcag	cctgtggggc	aggacggggg	1200
ccttgccacc	tgcagcctga	ctgcgcgcga	agaccgggat	ccggtgggaa	tggaacctta	1260
cagctggagt	cgagagtctg	gggcaatggg	gcaggaaaag	tggtactaca	ttaccatctt	1320
tgctctcgcg	caccccgaga	agctcacctt	gtggtctacg	gtcctgtcca	cctaccactt	1380
tgggggcaat	gcctcagcag	ctgggacacc	gcaccacgtc	tcggtgaaga	atcatagctt	1440
ggactctgtg	tctgtggact	gggcaccatc	cctgctgagc	acctgtcccc	gcgtcctaaa	1500
ggagtatgtt	gtccgctgcc	gagatgaaga	cagcaaacag	gtgtcagagc	atcccgctga	1560
gcccacagag	acccaagtta	ccctcagtgg	cctgcgggct	ggtgtagcct	acacggtgca	1620
ggtgcgagca	gacacagcgt	ggctgagggg	tgtctggagc	cagccccagc	gcttcagcat	1680
cgaagtgcag	gtttctgatt	ggctcatctt	cttcgcctcc	ctggggagct	tcttgagcat	1740
ccttctcgtg	ggcgtccttg	gctaccttgg	cctgaacagg	gccgcacggc	acctgtgccc	1800
gccgtgcccc	acaccctgtg	ccagctccgc	cattgagttc	cctggaggga	aggagacttg	1860
gcagtggatc	aaccagctgg	acttcagga	agaggcatcc	ctgcaggagg	ccctggtggt	1920
agagatgtcc	tgggacaaag	gcgagaggac	tgagcctctc	gagaagacag	agctacctga	1980
gggtgccccct	gagctggccc	tgatacaga	gttgtccttg	gaggatggag	acaggtgcaa	2040
ggccaagatg	tgatcgttga	ggctcagaga	gggtgagtga	ctgccccgag	gctacgtagc	2100

<210> SEQ ID NO 151

<211> LENGTH: 662

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 151

Met Glu Pro Leu Val Thr Trp Val Val Pro Leu Leu Phe Leu Phe Leu
 1 5 10 15

-continued

Leu Ser Arg Gln Gly Ala Ala Cys Arg Thr Ser Glu Cys Cys Phe Gln
 20 25 30
 Asp Pro Pro Tyr Pro Asp Ala Asp Ser Gly Ser Ala Ser Gly Pro Arg
 35 40 45
 Asp Leu Arg Cys Tyr Arg Ile Ser Ser Asp Arg Tyr Glu Cys Ser Trp
 50 55 60
 Gln Tyr Glu Gly Pro Thr Ala Gly Val Ser His Phe Leu Arg Cys Cys
 65 70 75 80
 Leu Ser Ser Gly Arg Cys Cys Tyr Phe Ala Ala Gly Ser Ala Thr Arg
 85 90 95
 Leu Gln Phe Ser Asp Gln Ala Gly Val Ser Val Leu Tyr Thr Val Thr
 100 105 110
 Leu Trp Val Glu Ser Trp Ala Arg Asn Gln Thr Glu Lys Ser Pro Glu
 115 120 125
 Val Thr Leu Gln Leu Tyr Asn Ser Val Lys Tyr Glu Pro Pro Leu Gly
 130 135 140
 Asp Ile Lys Val Ser Lys Leu Ala Gly Gln Leu Arg Met Glu Trp Glu
 145 150 155 160
 Thr Pro Asp Asn Gln Val Gly Ala Glu Val Gln Phe Arg His Arg Thr
 165 170 175
 Pro Ser Ser Pro Trp Lys Leu Gly Asp Cys Gly Pro Gln Asp Asp Asp
 180 185 190
 Thr Glu Ser Cys Leu Cys Pro Leu Glu Met Asn Val Ala Gln Glu Phe
 195 200 205
 Gln Leu Arg Arg Arg Gln Leu Gly Ser Gln Gly Ser Ser Trp Ser Lys
 210 215 220
 Trp Ser Ser Pro Val Cys Val Pro Pro Glu Asn Pro Pro Gln Pro Gln
 225 230 235 240
 Val Arg Phe Ser Val Glu Gln Leu Gly Gln Asp Gly Arg Arg Arg Leu
 245 250 255
 Thr Leu Lys Glu Gln Pro Thr Gln Leu Glu Leu Pro Glu Gly Cys Gln
 260 265 270
 Gly Leu Ala Pro Gly Thr Glu Val Thr Tyr Arg Leu Gln Leu His Met
 275 280 285
 Leu Ser Cys Pro Cys Lys Ala Lys Ala Thr Arg Thr Leu His Leu Gly
 290 295 300
 Lys Met Pro Tyr Leu Ser Gly Ala Ala Tyr Asn Val Ala Val Ile Ser
 305 310 315 320
 Ser Asn Gln Phe Gly Pro Gly Leu Asn Gln Thr Trp His Ile Pro Ala
 325 330 335
 Asp Thr His Thr Glu Pro Val Ala Leu Asn Ile Ser Val Gly Thr Asn
 340 345 350
 Gly Thr Thr Met Tyr Trp Pro Ala Arg Ala Gln Ser Met Thr Tyr Cys
 355 360 365
 Ile Glu Trp Gln Pro Val Gly Gln Asp Gly Gly Leu Ala Thr Cys Ser
 370 375 380
 Leu Thr Ala Pro Gln Asp Pro Asp Pro Ala Gly Met Ala Thr Tyr Ser
 385 390 395 400
 Trp Ser Arg Glu Ser Gly Ala Met Gly Gln Glu Lys Cys Tyr Tyr Ile
 405 410 415
 Thr Ile Phe Ala Ser Ala His Pro Glu Lys Leu Thr Leu Trp Ser Thr
 420 425 430

-continued

Val Leu Ser Thr Tyr His Phe Gly Gly Asn Ala Ser Ala Ala Gly Thr
435 440 445

Pro His His Val Ser Val Lys Asn His Ser Leu Asp Ser Val Ser Val
450 455 460

Asp Trp Ala Pro Ser Leu Leu Ser Thr Cys Pro Gly Val Leu Lys Glu
465 470 475 480

Tyr Val Val Arg Cys Arg Asp Glu Asp Ser Lys Gln Val Ser Glu His
485 490 495

Pro Val Gln Pro Thr Glu Thr Gln Val Thr Leu Ser Gly Leu Arg Ala
500 505 510

Gly Val Ala Tyr Thr Val Gln Val Arg Ala Asp Thr Ala Trp Leu Arg
515 520 525

Gly Val Trp Ser Gln Pro Gln Arg Phe Ser Ile Glu Val Gln Val Ser
530 535 540

Asp Trp Leu Ile Phe Phe Ala Ser Leu Gly Ser Phe Leu Ser Ile Leu
545 550 555 560

Leu Val Gly Val Leu Gly Tyr Leu Gly Leu Asn Arg Ala Ala Arg His
565 570 575

Leu Cys Pro Pro Leu Pro Thr Pro Cys Ala Ser Ser Ala Ile Glu Phe
580 585 590

Pro Gly Gly Lys Glu Thr Trp Gln Trp Ile Asn Pro Val Asp Phe Gln
595 600 605

Glu Glu Ala Ser Leu Gln Glu Ala Leu Val Val Glu Met Ser Trp Asp
610 615 620

Lys Gly Glu Arg Thr Glu Pro Leu Glu Lys Thr Glu Leu Pro Glu Gly
625 630 635 640

Ala Pro Glu Leu Ala Leu Asp Thr Glu Leu Ser Leu Glu Asp Gly Asp
645 650 655

Arg Cys Lys Ala Lys Met
660

<210> SEQ ID NO 152

<211> LENGTH: 360

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 152

caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc	60
acctgcactg tctctggtgg ctccatcagc agtgggtggtt actactggag ctggatccgc	120
cagcaccag ggaagggcct ggagtggatt gggcacatcc attacagtgg gaacacctac	180
tacaaccgct ccctcaagag tcgagttacc atatcagtag acacgtctaa gaatcagttc	240
tccctgaaac tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgcgaaat	300
cgcggttct actacggtat ggacgtctgg ggccaaggga ccacggtcac cgtctcctca	360

<210> SEQ ID NO 153

<211> LENGTH: 120

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 153

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln	
1 5 10 15	
Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly	
20 25 30	

-continued

Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu
 35 40 45

Trp Ile Gly His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser
 50 55 60

Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe
 65 70 75 80

Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr
 85 90 95

Cys Ala Arg Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 154
 <211> LENGTH: 23
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Honeybee melittin signal

<400> SEQUENCE: 154

Met Lys Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile
 1 5 10 15

Ser Tyr Ile Tyr Ala Ala Ala
 20

<210> SEQ ID NO 155
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: His Tag

<400> SEQUENCE: 155

His His His His His His
 1 5

What is claimed is:

1. An isolated nucleic acid molecule encoding one or both variable regions of an antigen binding protein that binds IL-23, wherein the antigen binding protein comprises at least one heavy chain variable region comprising: a CDRH1 of SEQ ID NO: 91, a CDRH2 of SEQ ID NO:92, and a CDRH3 of SEQ ID NO: 93; and at least one light chain variable region comprising: a CDRL1 of SEQ ID NO: 62, a CDRL2 of SEQ ID NO:63, and a CDRL3 of SEQ ID NO:64.
2. An isolated nucleic acid molecule encoding one or both variable regions of an isolated antigen binding protein that binds IL-23, wherein the antigen binding protein comprises a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-113 of SEQ ID NO:31; and a light chain variable region comprising amino acid residues 23-36, 52-58 and 91-101 of SEQ ID NO:1.
3. An isolated nucleic acid molecule encoding one or both variable regions of an isolated antigen binding protein that binds IL-23, wherein the antigen binding protein comprises a heavy chain variable region of SEQ ID NO: 31 and a light chain variable region of SEQ ID NO: 1.
4. The isolated nucleic acid molecule of claim 3, wherein the nucleic acid encoding the heavy chain variable region is SEQ ID NO:32.
5. The isolated nucleic acid molecule of claim 3, wherein the nucleic acid encoding the light chain variable region is SEQ ID NO:2.

6. The isolated nucleic acid molecule of claim 3, wherein the nucleic acid encoding the heavy chain variable region is SEQ ID NO:32 and the nucleic acid encoding the light chain variable region is SEQ ID NO:2.

7. A nucleic acid molecule according to claim 1, 2 or 3, wherein said nucleic acid molecule is operably linked to a control sequence.

8. A nucleic acid molecule according to claim 6, wherein said nucleic acid molecule is operably linked to a control sequence.

9. An isolated vector comprising a nucleic acid molecule according to claim 1, 2 or 3.

10. An isolated vector comprising a nucleic acid molecule according to claim 6.

11. An isolated host cell comprising the nucleic acid molecule according to claim 1, 2 or 3.

12. An isolated host cell comprising the nucleic acid molecule according to claim 6.

13. An isolated host cell comprising the vector according to claim 9.

14. An isolated host cell comprising the vector according to claim 10.

15. A method of making an antigen binding protein that binds IL-23 comprising the step of collecting said antigen binding protein from a host cell of claim 11, wherein the host

165

cell comprises nucleic acid encoding both the heavy chain variable region and the light chain variable region of the antigen binding protein.

16. A method of making an antigen binding protein that binds IL-23 comprising the step of collecting said antigen binding protein from a host cell of claim 12, wherein the host cell comprises nucleic acid encoding both the heavy chain variable region and the light chain variable region of the antigen binding protein. 5

17. A method of making an antigen binding protein that binds IL-23 comprising the step of collecting said antigen binding protein from a host cell of claim 13, wherein the host cell comprises nucleic acid encoding both the heavy chain variable region and the light chain variable region of the antigen binding protein. 10 15

18. A method of making an antigen binding protein that binds IL-23 comprising the step of collecting said antigen binding protein from a host cell of claim 14, wherein the host cell comprises nucleic acid encoding both the heavy chain variable region and the light chain variable region of the antigen binding protein. 20

* * * * *

166